

Health Effects of Nitrogen in Drinking Water

Mary H. Ward, Ph.D.

Text

Occupational and Environmental Epidemiology Branch
Division of Cancer Epidemiology and Genetics
National Cancer Institute, National Institutes of Health

Reducing the Health Impacts of the Nitrogen Problem: Defining the Problem (Workshop Session #1)



**NATIONAL
CANCER
INSTITUTE**


The National Academies of
SCIENCES • ENGINEERING • MEDICINE

Overview

- **Biologic effects of ingestion of nitrate in drinking water**
 - ❖ **N-nitroso compound (NOC) formation**
- **Health effects**
 - ❖ **Methemoglobinemia**
 - ❖ **Adverse reproductive outcomes**
 - ❖ **Cancers**
 - ❖ **Thyroid disease**

Nitrate in drinking water: Sources and exposures

- Nitrogen fertilizers, animal and human waste
- Regulatory limit (Maximum Contaminant Level [MCL]):
 - 10 mg/L as $\text{NO}_3\text{-N}$ (USA)
 - 50 mg/L as NO_3 (EU)
- Highest exposures:
 - Residents of agricultural areas
 - Private wells
 - Not regulated
 - Sparse measurements





Nolan et al, EST 2002

Nitrate in the diet: Sources and exposure

- High nitrate vegetables (e.g., green leafy vegetables, celery, beets)
- Vegetables also contain vitamin C, polyphenols, & other antioxidants inhibits formation of N-nitroso compounds (NOC)
- Beneficial effects on the cardiovascular system through nitric oxide (NO) formation
- Supplements with high nitrate (e.g. beet juice) have not been evaluated for NOC formation
- Nitrate & nitrite added to processed/cured meats linked to colorectal and other cancers



Nitrate in Vegetables (mg/100 g)



N-nitroso compound (NOC) formation from ingested nitrate (drinking water & diet)

Oral bacteria: Nitrate → nitrite


Nitrite + stomach acid

N_2O_3 +
amines/amides

NOC


Heme iron
(red meat)
Thiocyanate
(smoking)

Antioxidants
(vitamin C)




Human studies of nitrate ingestion

- 5-8% of ingested nitrate reduced to nitrite by the oral bacteria
- Direct relationship between nitrate conversion to nitrite and NOC concentrations in urine
- Drinking water nitrate increases endogenous formation of NOC:
 - >10 mg/L $\text{NO}_3\text{-N}$ (Mirvish 1992; Moller et al. 1989)
 - But not at levels <5 mg/L (Levallois, 2000)



Shapiro et al, Fd Chem Tox 1991

N-nitroso compounds (nitrosamines, nitrosamides): Cancer and birth defects



- ~300 NOCs tested in animals: 90% were carcinogenic (International Agency Research on Cancer [IARC], 1995)
- Carcinogens in 39 animal species including nonhuman primates
- Cause tumors in multiple organs: e.g., lung, digestive tract, bladder, kidney, ovary, thyroid
- *In utero* exposure causes congenital malformations especially of the central nervous system
 - ❖ First trimester most important for organ development

Methemoglobinemia

Nitrate and infant methemoglobinemia

September 8, 1945

CYANOSIS IN INFANTS CAUSED BY NITRATES IN WELL WATER

HUNTER H. COMLY, M.D.

» Author Affiliations

JAMA. 1945;129(2):112-116. doi:10.1001/jama.1945.02860360014004



- **Nitrite binds to hemoglobin forming methemoglobin (MetHb) and interferes with oxygen transport when >20% MetHb**
 - Infants <6 months most susceptible
- **Investigation of 279 cases in 14 U.S. states found no cases below 20 mg/L NO₃-N (Walton, Am J Public Health 1951)**
- **Case in WI at 27 mg/L NO₃-N (Knobeloch et al. Env Health Perspect 2000)**
- **Ongoing public health problem in Eastern Europe, The Gaza Strip, Morocco**

Adverse reproductive outcomes

- Spontaneous abortions
- Congenital malformations (birth defects)
- Few studies of other outcomes (low birth weight, preterm birth)

Adverse reproductive outcomes – spontaneous abortions

- High maternal MetHb can cause abortions in lab animals and livestock
- CDC MMWR report (1996) of a cluster of spontaneous abortions (miscarriage) in rural Indiana:
 - Women on private wells >20 mg/L NO₃-N
 - Switched to low nitrate water and had healthy live births
- Study in Massachusetts: no association with levels that were below 5.5 mg/L NO₃-N (Aschengrau, 1989)



Adverse reproductive outcomes:

Central Nervous System (CNS) malformations

- **Dorsch, 1984 (Australia)**: 3.5 increase in CNS defects for groundwater nitrate (<10 ppm nitrate-N)
- **Arbuckle, 1988 (Canada)**: two-fold increase in CNS malformations with nitrate >5.5 ppm in private wells
- **Croen, 2001 (California)**: increased risk anencephaly for nitrate >10 mg/L
- **Brender, 2004/2005 (Texas)**: two-fold increase neural tube defects (NTDs), nitrate >3.5 mg/L
- **Brender et al, 2013 (Texas, Iowa)**: two-fold increase in specific defects including spina bifida, limb deficiencies, cleft lip/palate
- **Holtby, 2014 (Canada)**: no association nitrate <MCL and all defects combined

Adverse reproductive outcomes: CNS malformations



- **Brender, 2004/2005 (Texas)**: two-fold increase in neural tube defects (NTDs)
 - Highest risk for women with high nitrate in their drinking water and who took prescription or over-the-counter drugs that can form NOC (nitrosatable drugs)
- **Brender, 2013 (Texas, Iowa)**: 2-fold increase in various defects: spina bifida, limb deficiencies, cleft lip/palate
 - Highest risk for women with high total nitrate intake (diet and drinking water) and nitrosatable drug use during pregnancy


**Summary: 5 of 6 studies found positive associations
nitrate concentrations were <10 mg/L in 4 studies**

Cancer

Ingested Nitrate and Nitrite (IARC vol. 94):

2A - probably carcinogenic to humans when ingested under conditions favorable for endogenous nitrosation

- Animal studies
 - 90% of NOC are carcinogens
 - Carcinogenic in all species tested
 - Many cancer sites
- Human mechanistic studies
 - NOC exposure increases with nitrate ingestion
- Epidemiology studies
 - Dietary nitrate – no studies with increased risk
 - Drinking water nitrate- inadequate evidence:
 - Few studies of specific cancer types
 - Low exposures - population using public water supplies
 - Historical exposures not always evaluated
 - Factors affecting endogenous nitrosation not always considered



IARC, 2010

Case-control or cohort studies of drinking water nitrate and cancer

Cancer: # studies (# positive in at least one subgroup):

- **Colorectal:** 5 studies (4 positive)
- **Bladder:** 4 studies (2 positive)
- **Kidney:** 2 studies (2 positive)
- **Brain (children):** 2 studies (2 positive)
- **Non-Hodgkin lymphoma:** 3 studies (1 positive)
- **Pancreas:** 2 studies (0 positive)
- **Brain (adults):** 2 studies (0 positive)

Iowa Women Health Study:




-42,000 postmenopausal women across Iowa

-73% use public water supplies, 25% private wells

**Public Water
Supply Monitoring
Data, 1955-1988**

NO₃-N

**Total Trihalomethanes
(THMs)**



Exposure Metrics

**Average NO₃-N and
THMs**

**Years >1/2 Maximum
Contaminant Level**

82% used supply 16+ years

Drinking water nitrate and cancer: early results



- **Weyer *et. al.* (2001)**

Ovarian cancer (n = 109)

- ❖ Public water: $HR_{Q4vs.Q1} = 2.0 (1.01 - 4.1)$
- ❖ Private well users (vs Q1): $HR = 1.6 (0.8 - 3.1)$

Bladder cancer (n=57)

- ❖ Public water: $HR_{Q4vs.Q1} = 2.8 (1.1 - 7.2)$
- ❖ Private well users (vs Q1): $HR = 1.3 (0.5 - 3.5)$


- **No association for colon, rectum, lung, breast, kidney, pancreas, NHL, melanoma**

Updated results: Ovary and Bladder Cancers





- **>20 years of follow-up:**
 - 315 ovary, 263 bladder cases
 - Similar increased risk (Inoue-Choi, 2015; Jones, 2016)
 - Updated analyses of colon, rectum, pancreas cancers (Jones, 2019; Quist, 2018)
 - Increased risk of kidney cancer in highest exposure group (Jones, 2017)
- **Exposure assessment:**
 - Trihalomethanes ($\rho=0.24$ with nitrate)
 - Dietary nitrite
- **Factors affecting NOC formation:**
 - Vitamin C and red meat
 - Smoking (bladder)

Average nitrate (mg/L) and ovarian cancer



Average nitrate and ovarian cancer, by vitamin C intake



$p\text{-interaction} = 0.01$

Summary and limitations of studies of drinking water nitrate and cancer




- **Most consistent evidence for colorectal cancer**
- **Still few studies of specific cancer sites**
- **Limitations of most studies:**
 - Private well users (highest exposures) usually not considered due to lack of measurement data
 - Factors affecting NOC formation not always evaluated

Ward et al. Int J Environ Res Pub Health, 2018

Thyroid disease

Nitrate inhibition of iodide uptake by the thyroid

Thyroid and Parathyroid Glands



Nitrate in drinking water and hypothyroidism

1. Subclinical hypothyroidism (measured by ultrasound) was increased in children in Slovakia with high nitrate drinking water (*Tajtakova et al, Chemosphere 2006*)
2. Postmenopausal women in Iowa with higher nitrate ingestion (diet and drinking water) had higher prevalence of hypothyroidism (*Ward et al, Epidemiology 2010*)
3. Amish women (but not men) in Pennsylvania with drinking water nitrate $>6 \text{ mg/L NO}_3\text{-N}$ had higher serum TSH & subclinical hypothyroidism (*Aschebrook-Kilfoy et al, Env Health 2012*)

Ongoing research on drinking water contaminants - the Agricultural Health Study cohort



- **Farmers and their spouses in Iowa and North Carolina (N=~90,000)**
- **60% use private wells at enrollment in 1993-1997**
- **Estimated nitrate in private wells (random forest models)**
- **Iowa private wells predicted nitrate: median (IQR): 1.6 (0.9-5.1), 13% > 10 mg/L**
Wheeler et al., STOTEN 2016
- **Public water supply measurements- nitrate, disinfection by-products**
- **Dietary nitrate/nitrite from food frequency questionnaire**

Research needs

- **Clarify relationship between nitrate ingestion & NOC formation below MCL in controlled studies**
- **Evaluate additional factors that influence endogenous nitrosation**
 - E.g. oral microbiome can be measured
 - Incorporate biomarkers (urinary NOC, nitrate) in studies with biological samples
- **Consider other water contaminants**
- **More studies in populations, high exposures**

Collaborators

National Cancer Institute

Brisa Aschebrook-Kilfoy

Kathryn Barry

Laura Beane Freeman

Kenneth Cantor

Curt DellaValle

Abby Flory

Maki Inoue-Choi

Rena Jones

U.S. Geologic Survey

Tom Nolan

University of Iowa

Peter Weyer

University of Minnesota

Kristin Anderson, DeAnn Lasovich, Kim Robien,
Anna Prizment

Mayo College of Medicine

James Cerhan

Virginia Commonwealth University

David Wheeler

Colorado State University

Jay Nuckols

Reviews of nitrate in drinking water and health

Workgroup Report: Drinking-Water Nitrate and Health—Recent Findings and Research Needs

VOLUME 113 | NUMBER 11 | November 2005

Mary H. Ward,¹ Theo M. deKok,² Patrick Levallois,³ Jean Brender,⁴ Gabriel Gulis,⁵ Bernard T. Nolan,⁶ and James VanDerslice⁷





International Journal of
*Environmental Research
and Public Health*



Review

Drinking Water Nitrate and Human Health: An Updated Review

Int. J. Environ. Res. Public Health **2018**, *15*, 1557

Mary H. Ward ^{1,*}, Rena R. Jones ¹ , Jean D. Brender ², Theo M. de Kok ³, Peter J. Weyer ⁴, Bernard T. Nolan ⁵, Cristina M. Villanueva ^{6,7,8,9}  and Simone G. van Breda ³