# Local tsunami warning for Cascadia: The challenges ahead

Diego Melgar University of Oregon














Central




UNIVERSITY of WASHINGTON




#### **Neskowin Ghost Forest, OR**



























#### Local Warning in Practice: The Tohoku Experience










Ozaki et al, 2011, EPS

#### 




Rapid determination of an earthquakes size and rupture characteristic is tricky.



## Fault Areas of large earthquakes




Rapid determination of an earthquakes size and rupture characteristic is tricky.



## So we use high precision GPS








## El GPS is the best sensor for very large EQs O | OREGON




#### 2011 M9 Tohoku-oki




#### We know it works





#### GPS also discriminates the "type" of earthquake O OREGON







## These happen in the real world











#### GPS in the western US






























Tsunami Amplitude

Tsunami Amplitude
















## How do we test the systems?



- Large earthquakes are infrequent, how do we know systems work?
- We create "scenarios" thousands of them and push them through the system.
- How well do we model earthquakes?
- How well do we model tsunamis?






Ruhl et al., GRL, 2017


#### Scenarios in Cascadia



- Large earthquakes are infrequent, how do we know systems work?
- We create
  "scenarios"
  thousands of them
  and push them
  through the system.
- How well do we model earthquakes?
- How well do we model tsunamis?







## So how are we doing?



- The outlook is overall positive, with continued effort warning for a Cascadia M9 earthquake in under 5 minutes will be possible.
- The key components, GPS and tsunami modeling, are there.
- We are now on the road from research to operations.
- Testing the operational environment will be important as well.
- --- Are we done?


## So how are we doing?



Earthquakes and tsunamis are complex


Washington

What I've described here is indirect warning, we measure the earthquake <u>not</u> the tsunami




What are the new technologies?





## Seismograph Network Operated by NIED (Inland)

- Real-time cabled sensors
- Will directly measure the tsunami
- The fastest most reliable way to issue accurate local warnings
- It is also useful for basic science
- It's costly, ~\$500M buildout and \$5M/yr operations



Strong motion seismograph network (K-NET&KiK-net)


- Measurement of strong ground motion.

Spacing of 20 km (1000+700stations)

Evaluation of source process for large earthquakes.

## A real-time ocean bottom network O oregon











## An example: The 2004 Indonesia Tsunami O | OREGON





## An example: The 2004 Indonesia Tsunami O | OREGON






## A tsunami's speed



15,000	500	Deep ocean
3000	222	Continental shelf
1500	158	Continental shelf
300	70	Coast


## When does the tsunami arrive?






#### When does the tsunami arrive?






## Second line of defense, buoys!






## How do we test the systems?






## Can we map out the shallow megatives region




## Can we map out the shallow megatives region



## The timeline for warnings





Tsunami warning centers have the operational goal to issue warnings in 20-30 minutes

#### A real-time test



M8.3 Tokachi-oki Japan earthquake

