Abundant spontaneous and dynamically triggered submarine landslides in the Gulf of Mexico

Wenyuan Fan


Assistant Professor
Institute of Geophysics and Planetary Physics
Scripps Institution of Oceanography
University of California, San Diego

In collaboration with Jeff McGuire, Peter Shearer, Catherine de Groot-hedlin, Michael Hedlin











Submarine landslides can damage offshore infrastructure






Taylor Energy site has been leaking since **2004** after the drilling platform was destroyed by a **submarine landslide**, the leaking would last for about **100 years** if there is no intervention




When, Where, thus Why?!


Abundant submarine landslides in the Gulf of Mexico





2008—2015, **85** in total Spontaneous: **10**Dynamically triggered: **75**




Direction of highly coherent local surface waves Vertical component (filtered at 20–50 s)














Centroid single force (CSF) inversion


















Centroid single force (CSF) inversion
















Centroid single force (CSF) inversion

~62 million tons of rocks/sediments






Submarine landslide dynamically triggered by a distant earthquake


Distance = 1547 kmTime = 435 s

Triggering velocity = 3.6 km/s



Submarine landslide dynamically triggered by a distant earthquake

Distance = 1547 kmTime = 435 sTriggering velocity = 3.6 km/s




This is **NOT** the near-field (< 100 km) type earthquake-landslide triggering process (PGA: ~1g?)

The earthquake and the submarine landslide are ~1500 km away.... (~KPa?)


75 dynamically triggered submarine landslides (2008–2015)

Detected over 250 triads


Detected over 150 triads




dynamically triggered submarine landslides (2008–2015)




Distance vs time of the triggered submarine landslides



Distance (km)







Detected over 150 triads

Detected over

250 triads

Fan et al., *GRL*, 2020







No magnitude dependence




Peak dynamic strain (stress) is not the only triggering threshold




Triggering magnitude histogram




Fan et al., *GRL*, 2020




Triggering distance histogram



Abundant submarine landslides in the Gulf of Mexico






2008—2015, **85** in total

Spontaneous: 10


Abundant submarine landslides in the Gulf of Mexico






2008—2015, **85** in total Spontaneous: **10**

Mechanisms of submarine landslides in the Gulf of Mexico



Complex ocean basin + thick sediment




fast, block, weak zone

2008—2015, **85** in total

Spontaneous: 10

Mechanisms of submarine landslides in the Gulf of Mexico




- Rapid sediment accumulation
- Overly steep topography
- Over pressurization
- Gas hydrate
- Weak oil layer
- Ground water / oil seepage
- Prolonged strong ground motion
- Cyclic shearing -> plastic strain -> strength reduction
- Cyclic shearing -> permeability enhancement

2008—2015, **85** in total

Spontaneous: 10


Mechanisms of submarine landslides in the Gulf of Mexico




- Complex ocean basin + thick sediment
- Rapid sediment accumulation
- Overly steep topography
- Over pressurization
- Gas hydrate
- Weak oil layer
- Ground water / oil seepage
- Prolonged strong ground motion
- Cyclic shearing -> plastic strain -> strength reduction
- Cyclic shearing -> permeability enhancement


2008—2015, **85** in total

Spontaneous: 10



Submarine Environmental Sensing Over Telecommunication Fiber Optic Cables





Cabled Observatories

Transoceanic Cables


Groundwater monitoring using ambient noise seismology

Marine Denolle, Tim Clements, Laura Ermert, Julian Schmitt



- Large-scale computing on the cloud
- 100-150 TB data archives
- 100s stations
- Julia scripting for fast processing

We could use ambient vibration to extrapolate or interpolate groundwater levels



Progress towards:

- Separating signals related to distinct environmental forces
- Monitoring and locating cliff failure
- Assessing influence of potential failure mechanisms

Outstanding challenges:

- Link between environmental forcing and cliff erosion rates
- Limited interrogation of influence and evolution of cliff material properties
- Limited focus on identifying precursors to cliff failure

Operational challenges and limitations:

- Small number of datasets w/ variable setups
- "Noisy" environments

- Directional sources
- Failure is rare, erosion hard to measure



Opportunities

- Identification of rare events, forcing/precursors and process cascades
- High spatial/temporal resolution for process monitoring
 - With independent calibration/data: direct estimates of sediment transport rates
 - Without independent calibration/data: tracking of qualitative patterns, relative rates
- Characterizing bed organization/evolution or energy transfer

Challenges/Needs

- Complex coupling between sediment transport, hydraulic parameters and environmental variables
- Site-specific attenuation (i.e., empirical Green's functions) riverbeds are heterogeneous!
- Parallel development and validation of theoretical models will require controlled seismic deployments co-located with traditional instrumentation to independently measure environmental variables
 - Weather/meteorological data
 - Water level/discharge
 - Sediment transport rates

- High resolution topography/change detection
- Substrate characteristics (attenuation)
- Video/time-lapse imagery

Innovative seismic techniques can be used to study the environment to benefit the society









