THE NATIONAL ACADEMIES OF SCIENCES, ENGINEERING, AND MEDICINE





The National Academy of Sciences

HE NATIONAL ACADEMY of Sciences was established in 1863 by an Act of Congress, signed by President Lincoln, as a private, nongovernmental institution to advise the nation on issues related to science and technology. Members are elected by their peers for outstanding contributions to research. Dr. Marcia K. McNutt is president.











The National Academy of Engineering was established in 1964 under the charter of the National Academy of Sciences to bring the practices of engineering to advising the nation. Members are elected by their peers for extraordinary contributions to engineering. Dr. C. D. Mote, Jr., is president.

The National Academy of Medicine (formerly the Institute of Medicine) was established in 1970 under the charter of the National Academy of Sciences to advise the nation on medical and health issues. Members are elected by their peers for dis-

tinguished contributions to medicine and health. Dr. Victor J. Dzau is president.

The three Academies work together as the National Academies of Sciences, Engineering, and Medicine to provide independent, objective analysis and advice to the nation and conduct other activities to solve complex problems and inform public policy decisions. The National Academies also encourage education and research, recognize outstanding contributions to knowledge, and increase public understanding in matters of science, engineering, and medicine.

W W W . N A T I O N A L - A C A D E M I E S . O R G

Dear Lecture Participant:

On behalf of the Ocean Studies Board at the National Academies of Sciences, Engineering, and Medicine, we would like to welcome you to the Nineteenth Annual Roger Revelle Commemorative Lecture. This lecture was created by the Ocean Studies Board in honor of Dr. Roger Revelle to highlight the important links between the ocean sciences and public policy.

ROGER REVELLE

For almost half a century, Roger Revelle was a leader in the field of oceanography.

Revelle trained as a geologist at Pomona College and the University of California,

Berkeley. In 1936, he received his Ph.D. in oceanography from the University of California, Berkeley. As a young naval officer, he helped persuade the Navy to create the Office of Naval Research (ONR) to support basic research in oceanography and was the first head of ONR's geophysics branch. Revelle served for 12 years as the Director of Scripps (1950–1961, 1963–1964), where he built up a fleet of research ships and initiated a decade of expeditions to the deep Pacific that challenged existing geological theory.

Revelle's early work on the carbon cycle suggested that the sea could not absorb all the carbon dioxide released from burning fossil fuels. He organized the first continual measurement of atmospheric carbon dioxide, an effort led by Charles Keeling, resulting in a long-term record that has been essential to current research on global climate change. With Hans Suess, he published the seminal paper demonstrating the connection between in-

creasing atmospheric carbon dioxide and burning of fossil fuels. Revelle kept the issue of increasing carbon dioxide levels before the public and spearheaded efforts to investigate the mechanisms and consequences of climate change. Revelle left Scripps for critical posts as Science Advisor to the Department of the Interior (1961–1963) and as the first Director of the Center for Population Studies at Harvard (1964–1976). Revelle applied his knowledge of geophysics, ocean resources, and population dynamics to the world's most vexing problems: poverty, malnutrition, security, and education.

In 1957, Revelle became a member of the National Academy of Sciences to which he devoted many hours of volunteer service. He served as a member of the Ocean Studies Board, the Board

2018 OCEAN STUDIES BOARD MEMBERS Larry A. Mayer, Chair, University of New Hampshire, Durham | E. Virginia Armbrust, University of Washington, Seattle | Kevin R. Arrigo, Stanford University, California | Claudia Benitez-Nelson, University of South Carolina, Columbia | Thomas S. Chance, ASV Global, LLC, Broussard, Louisiana | Rita R. Colwell, University of Maryland, College Park | Sarah W. Cooksey, Delaware Chapter of the Nature Conservancy, Milton | James A. Estes, University of California, Santa Cruz | David Halpern, Jet Propulsion Laboratory, Pasadena, California | Patrick Heimbach, University of Texas, Austin | Susan E. Humphris, Woods Hole Oceanographic Institution, Woods Hole, Massachusetts | S. Bradley Moran, University of Alaska, Fairbanks | Steven A. Murawski, University of South Florida, St. Petersburg | John A. Orcutt, Scripps Institution of

on Atmospheric Sciences and Climate, and many committees. He also chaired a number of influential Academy studies on subjects ranging from the environmental effects of radiation to understanding sea-level change.

SMITHSONIAN'S NATIONAL MUSEUM OF NATURAL HISTORY

The Ocean Studies Board is pleased to have the opportunity to present the Revelle Lecture in cooperation with the Smithsonian National Museum of Natural History through our partnership with the Smithsonian Science Education Center. The museum maintains and preserves the world's most extensive collection of natural history specimens and human artifacts and supports scientific research, educational programs, and exhibitions. The museum is part of the Smithsonian Institution, the world's largest museum and research complex. Dr. Kirk R. Johnson is the director.

The Smithsonian Science Education Center (SSEC) was founded in 1985 by the National Academy of Sciences and the Smithsonian Institution and continues today as a successful unit of the Smithsonian Institution. The mission of the SSEC is to develop STEM literate students from early childhood through the workplace. The SSEC does this through the implementation of a truly systemic approach that engages participants at every level, from students and classroom teachers up through the highest levels of district, state, national and international leadership.

TONIGHT'S LECTURE

Many fish stocks have been reduced to a fraction of their former abundance, the result of overfishing, habitat destruction, pollution, and other human impacts. However, fisheries managers often set baselines for rebuilding to a time when fish stocks may already have been depleted. In his lecture this evening, Dr. Jeffrey Bolster, professor of history at the University of New Hampshire, uncovers the deep roots of the depletion of our coastal ecosystems. Reconstructing the catch histories of commercial fisheries in the northwest Atlantic in the 19th century, Dr. Bolster concludes that we may be profoundly underestimating the capacity of the ocean to produce fish.

SPONSORSHIP

The Ocean Studies Board thanks the National Oceanic and Atmospheric Administration, the National Science Foundation, the National Aeronautics and Space Administration, the Office of Naval Research, and the U.S. Geological Survey. This lecture series would not be possible without their generous support.


We hope you enjoy tonight's event.

CHAIR, OCEAN STUI

Susan Roberts.

DIRECTOR, OCEAN STUDIES BOARD

Oceanography, La Jolla, California | H. Tuba Özkan-Haller, Oregon State University, Corvallis | Ruth M. Perry, Shell Exploration & Production Company, Houston, Texas | Martin D. Smith, Duke University, Durham, North Carolina | Mark J. Spalding, The Ocean Foundation, Washington, District of Columbia | Margaret Spring, Monterey Bay Aquarium, Monterey, California | Douglas Wartzok, Florida International University, Miami | Lisa D. White, University of California, Berkeley and San Francisco State University | Robert S. Winokur, Michigan Tech Research Institute, Silver Spring, Maryland | OSB STAFF MEMBERS Susan Roberts, Director | Stacee Karras, Program Officer | Emily Twigg, Associate Program Officer | Pamela Lewis, Administrative Coordinator | Trent Cummings, Program Assistant | Shubha Banskota, Financial Associate | James Heiss, Post-Doctoral Fellow



PROFESSOR OF HISTORY at the University of New Hampshire, Jeff Bolster helped create the emerging field of marine environmental history. For ten years, he co-directed the UNH Cod Project, part of the Census of Marine Life's historical arm. Bolster's interdisciplinary group of ecologists and historians pioneered new methods to reconstruct the abundance and distribution of historic fish stocks in the northwest Atlantic. He continues to advocate for the importance of historical evidence in investigating scientific questions about long-term environmental change.

DR. W. JEFFREY BOLSTER







Bolster went to sea for ten years as a young man, sailing as mate and master on a variety of sailing school-ships, including the Sea Education Association's R/V Westward. Licensed by the U.S. Coast Guard as Master of Motor, Steam, and Auxiliary Sail Vessels upon All Oceans, he still regularly sails small vessels deep-sea, and remains intimately familiar with the ocean and the fisheries of New England, Atlantic Canada, and the eastern Caribbean.

He is author, co-author, or editor of five books, many papers, and contributions to the New York Times, the Boston Globe, and other publications. Bolster's first book, Black Jacks: African American Seamen in the Age of Sail won several academic prizes, and the New York Times Book Review listed it as a "notable book of the year." His most recent book, The Mortal Sea: Fishing the Atlantic in the Age of Sail, won numerous awards, including the

Bancroft Prize, generally regarded as one of the most prestigious awards in American History. A reviewer in The Washington Post said, "Anyone who thinks ... this book is only about fish is living in a fool's paradise."

Bolster has held fellowships from the Smithsonian Institution, Mystic Seaport, and the National Endowment for the Humanities, and he served one year as the Fulbright Distinguished Chair in American Studies at the University of Southern Denmark. His work has been supported by grants from the Alfred P. Sloan Foundation, the National Science Foundation, New Hampshire Sea Grant, and NOAA's Marine Sanctuary Program.

Educated at Trinity College (BA, History), Brown University (MA, History), and the Johns Hopkins University (PhD, History), Bolster lives in Portsmouth, New Hampshire, an old town by the sea.

Y INVITING A HISTORIAN to deliver the 2018 Roger Revelle Lecture, the Ocean Studies Board signals the value of fruitful engagement between the sciences and humanities. Their decision honors Roger Revelle's legacy. A brilliant geophysicist and ocean-

ographer, Revelle embraced interdisciplinary approaches during the final years of his career, turning his attention to malnutrition, global poverty, and education. Roger Revelle recognized that thorny problems required multiple forms of knowledge. Historians approach the past as if it were a foreign country. Getting there is difficult; understanding what's encountered even more so. (Lowenthal, 1985; Appleby et al., 1994) Evidence is fragmentary, and contexts are crucial. Historical observations are clearly important: anyone seeking perspective on the contemporary fisheries crisis, for instance, or striving to understand how healthy marine ecosystems once functioned, would do well to pay attention to the past. Yet despite a recent groundswell of interest in historical evidence by some marine scientists, it has yet to penetrate deeply into assessment and policy (Alexander et al., 2011; Kittinger et al., 2015; Engelhard et al., 2016).











We are the first generation to confront numerous distress signals from the living ocean: pollution, habitat destruction, plastic infestation, anoxic zones, overfishing, biological invasions, ocean acidification, and coral bleaching. Worries in the past were more focused. In nineteenthcentury America, they revolved around a central question: "Would there be fish for the future?"

It is easy to blame modern technology for ecological problems in the fishery, easy to assume that our problems began with post-World War Two factory ships, rugged polyester nets, electronic fishfinders and pinpoint GPS navigation. Voluminous evidence shows otherwise. Overfishing has deep

roots in the northwest Atlantic, and in much of the world (Jackson et al., 2001; Myers and Worm, 2003; Rosenberg et al., 2005; Lotze et al, 2006; Bolster, 2012). When historical evidence is assessed, one thing becomes readily apparent. The state of marine ecosystems and fisheries is worse than most experts imagine.

American fishermen, scientists,



Figure 1. During the 1850s when serious concerns were first raised about the depletion of cod and mackerel stocks, this is the sort of simple vessel from which fish were caught. Men hand-lined from deck. The Marblehead schooner, Amy Knight, built c. 1820. Model by Erik Ronnberg. Photo courtesy of Erik Ronnberg.

and policymakers have been grappling with fisheries depletion for a very long time. That conversation began in earnest during the 1850s (Figure 1) (Bolster, 2012). In 1871 federal politicians commenced spending taxpayers' dollars to reverse declining catches. Spencer F. Baird, one of the most prominent scientists in America, and the Director of the U.S. Commission on Fish and Fisheries, argued shortly thereafter for "the restoration of our exhausted cod fisheries" (USCFF, 1874). It never happened (Figure 2).

By the late nineteenth and early twentieth centuries, the U.S. Commission on Fish and Fisheries, and its successor after 1903, the U.S. Bureau of Fisheries, oversaw one of the most generously funded federal research initiatives in the country (Pauly, 2000; Bolster, Maclean, 2003; Rose, 2007). 2012). Investigators' data revealed on-going depletion.

Industrial fishing began in earnest after World War One. Overfishing increased. By the 1950s, foreign factory processing ships were fishing the northwest Atlantic, further reducing biomass (Warner, 1977). By 1976, with the adoption of the Magnuson Act and the 200-mile Exclusive Economic Zone, northwest Atlantic groundfish stocks were reeling. They have never recovered. Many have continued to decline, despite

scientific management under revisions of federal fisheries law in the Magnuson-Stevens Act (Pauly and

Our twenty-first-century fisheries crisis has been regarded rightfully as an ecological and political problem, but too infrequently understood in light of history—as if nature and science were somehow separate from the study of the past (Bolster, 2006). The lion's share of marine ecology and fisheries management articles published every year cover only a brief span of time. They cannot help but suspend attention to drivers of ecosystem function such as natural variability, historic anthropogenic

influences, the presence or absence of certain organisms, long-term climatic cycles, and extreme weather events, among others. Lack of data is clearly a problem, compounded by some researchers' unwillingness to dig deeply in unfamiliar types of sources, or to imagine how various materials from the past might inform their studies.

Statistical stock assessments remain the gold standard for management decisions, yet they rarely rely on information more than a few decades old. Managers' goal is sustaining the biomass necessary for maximum sustainable yield. Recent data alone, however, and calculations derived from it (no matter how elegant), cannot help but ignore historic ecosystem productivity, along with other aspects of past ecosystem configuration. That leads policy-makers, pressured by commercial and community groups, to establish restoration goals reflecting only a fraction of fish abundances once considered normal. Such present-centered approaches lead everyone—scientists, policy-makers, and the public—to misapprehend the scale of the problem, and the fact that generations have struggled with it.

I admit candidly that historical evidence is often messy, incomplete, and inconsistent—that it is characterized by uncertainties. Frequently it does not lend itself to quantification. It is often difficult to merge with data recently obtained.

Of course, sophisticated stock assessments and mathematically

precise ecosystem modeling are also riddled with uncertainties, depending on the assumptions made, data used, categories of analysis ignored, and hypotheses examined (Link et al., 2012). Each approach, whether by historians or modelers, has uncertainties. They are just different, a result of those practitioners' training and the material at their disposal. The anthropologist



FIGURE 2. In 1873, Spencer F. Baird, Director of the U.S. Fish Commission, argued for "restoration of our exhausted cod fisheries." It never happened. Photo courtesy of NOAA. Adapted from W. Jeffrey Bolster, The Mortal Sea: Fishing the Atlantic in the Age of Sail (2012).

Ruth Benedict pointed out years ago that if we are to truly understand things, "we must know as much about the eye that sees as about the object seen Means of perception [are] conditioned by the trajectory in which its possessor has been reared" (Benedict, 1943). Fisheries management, I suggest, would benefit from openness to more "means of perception."

Can we expand the definition of what constitutes the "best scientific information available" to include more evidence from the past? A National Academies report in 2004 on improving the best scientific information available took a step in that direction when it referred to the usefulness of anecdotal information in some circumstances. Of course, anecdotes are only one form of historical evidence (NRC, 2004).

The conversation is really about that old question—will there be fish for the future? Managers need tools to help stocks recover. It is easy to make a case that without statistical stock assessment and scientific management since 1976, the state of the fisheries would be considerably worse than it currently is. Pressures from the fishing industry and its lobbyists have been intense. That said, management protocols in place during the last forty years have not stopped the on-going degradation of many fisheries. In the face of this on-going problem, methodological change seems warranted.

Meaningful change to the federal fisheries law should include recognition of findings by historical marine ecologists. Historical observations provide sorely needed perspective on rebuilding fish stocks. Clearly, protocols for evaluating such work will need to be developed. While that will be contentious, management approaches honoring different forms of knowledge should contribute to the recovery of living marine resources.

I certainly don't have all the answers, but I can tell you how we got here.



ROM 1639 to 1702 New Englanders passed laws to preserve striped bass, cod, and mackerel,

despite the sea of plenty lapping at their feet. We could snort with derision, regarding those colonists as foolish because there were plenty of fish. Or we could recognize that as emigrants from places with degraded ecosystems, colonials were worried about depletion. Their policies reflected local ecological concerns (Bolster, 2012).

Discernable human impacts on marine ecosystems were apparent by the presidency of George Washington. By then, Atlantic Gray Whales were extinct. Right Whales were greatly reduced. Great Auks, North Atlantic "penguins," were heading for extinction. Other seabirds had been depleted. The abundance and distribution of walrus had been severely reduced, pushing a species once common near Nova Scotia and New Brunswick toward the Arctic. Anadromous fish runs had shrunk

In 1797 Judge Benjamin Chadbourne described the Salmon Falls River, which flows into the Piscataqua River, the southern border between Maine and New Hampshire. "Formerly, large fish such as

salmon, bass and shad came up the river in plenty, but they have forsook it." A keen observer and lifelong resident, Chadbourne, then in his sixties, revealed how fishing had altered the estuarine ecosystem during his lifetime. Evidence from the northwest Atlantic prior to 1800 reveals that the ecosystem roles of marine mammals, sea birds, and anadromous fish had all been reduced, and that estuarine productivity had been degraded. Such observations are waypoints worth noting (Chadbourne, 1797).

American fishermen ignored menhaden entirely throughout the colonial period, though farmers in Long Island Sound and Narragansett Bay seined them during the

"Each

human

came to

expect

settling

radically

natural

world"

less.

for a

late eighteenth century for fertilizer. Menhaden are bony, oily cousins of herring, and forage fish par excellence (Franklin, 2007). Today industrial fishers seine menhaden to render them into oil, and animal or poultry feed. A small-scale menhaden oil rendering operation began in Rhode Island in 1811, but fishermen in Maine generally continued to ignore those little fish.

That changed in 1850. John Bartlett, of Blue Hill, Maine seined a few baskets of menhaden, which his wife boiled on the beach. She skimmed the oil, and a Boston merchant offered her \$11 a barrel.

The Bartletts seined more ambi-

tiously—still with handmade nets from small rowboats and sailboats near shore. Neighbors got involved

People in Blue Hill had commercialized a previously underutilized resource—a recurring theme in fisheries history. Otherwise, they were doing with menhaden what they and neighbors had done with cod and mackerel for decades. Locals caught nearby fish, packed fish products in barrels, and sent them away for money. It seemed logical and lucrative (Bolster, 2012).

Yet hundreds of fishermen reacted angrily, inundating legislators with petitions. Boothbay residents in 1852 insisted that "Taking Menhaden ... in our Bays, Rivers, and Harbours is very destructive to said

fish and if persisted in will eventually destroy them or drive them from our coast." Men from Deer **generation** Isle, Ellsworth, Surry, and Sedgewick concurred. In 1857 Gouldsboro fishermen, fearing "the material injury of the codfishing interests in this state, requested legislative action diminished to "prevent the future destruction of the menhaden" (MeSA, Legislative Laws, 1852; MeSA, Legislative Graveyard, 1857).

> This fury was unlike anything that had previously existed in Maine's legislative record. Upon encountering it in the archives, I was shocked. Why were so many fishermen angry about the commercialization of menhaden, and

the money it was bringing into their hardscrabble towns?

As pieces of the puzzle came together, it became clear that the 1850s were the first decade in which fishermen from Massachusetts, New Hampshire, Maine, Nova Scotia, and New Brunswick organized systematically to express concerns about the future of cod, haddock, and mackerel. Colonials had lamented the demise of anadromous fish—the shad, salmon, sturgeon, and others that ascended rivers to spawn. Those fish were sitting ducks. Not until the 1850s, however, did large numbers of fishermen articulate what they saw as threats to true sea fish. In 1852 the Nova Scotia Assembly debated outlawing the mackerel fishery

during spawning season. In 1855 the Maine Legislature prohibited seining mackerel within three miles of shore—the area it controlled (Bolster, 2012). Historians call what occurred in the 1850s a turning point. By then, the fishing banks seaward of New England and Atlantic Canada had been a coupled human-and-natural system for centuries (Liu et al., 2007).





FIGURE 3. Geographic distribution of 19th-century fishing, with modern management areas. The Nova Scotian Shelf, highlighted in blue, figured prominently on this 1853 British Admiralty Chart of the Gulf and River St. Lawrence ... (London: Bayfield, Holbrook, & Bullock). Red crosses track the course of the Beverly schooner Angler, spring 1853. Red ovals indicate location and relative magnitude of daily catch. British Admiralty chart courtesy of Peabody Essex Museum. Chart of NAFO management areas courtesy Northwest Atlantic Fisheries Organization. GIS by Stefan Claesson. Compiled by Karen E. Alexander. Adapted from Rosenberg, et. al., 2005, Frontiers in Ecology and The Environment.



came together in 2000, committed to interdisciplinary approaches to historic fisheries. We were fortunate to find historical records that could be merged with a modern population dynamics model to create quantifiable data for nineteenth century fisheries. Our results helped explain why fishermen changed their tune during the 1850s. Fishermen felt that catches were declining. They were right.

One of our studies used extraordinarily rich data from 326 fishing schooners from Beverly, Massachusetts that fished the Nova Scotian Shelf during the 1850s, along with catch records from another 1,313 American schooners also fishing there (Rosenberg et al., 2005; Bolster et al., 2011).

We discovered that the 1850s were a grim decade in the offshore banks fishery (Figure 3). Landings per boat per season declined from 26,217 cod in 1852 to only 14,414 in 1859. Statisticians in our group analyzed the data using a variation of the Chapman-DeLury stock assessment method, assuming identical rates of natural mortality and recruitment. The model allowed solving for initial abundance, that is, the biomass of cod on the Nova Scotian Shelf in 1852. Our estimate was 1.26 million metric tons (mt) (Figure 4).

The 2002 cod biomass estimate by Fisheries and Oceans Canada for the same region was less than 50,000 metric tons, about 4% of that in 1852 (Canada DFO, 2002). Cod biomass there has continued to

decline. The 2009 biomass estimate was a paltry 24,000 mt, only 2% of that in 1852. Given very light fishing pressure in recent years, it appears that natural mortality is decimating cod (Canada DFO, 2017).

Numerical estimates of historic fisheries abundance are very rare. Few extend beyond the 1960s, even in the North Atlantic, where data collection has been the most comprehensive. Fisheries records from most other parts of the world are notoriously worse, characterized by short time spans, unreliable landings records, and lack of catchper-unit-effort (CPUE) data (Al-Abdulrazzak et al., 2015). Put another way, we don't know a great deal about measurable ecosystem productivity in many areas today, much less in the past. Creating a defensible biomass estimate for an important fishing ground in 1852, before the onset of industrialized fishing, provided the scale of a healthy fish population. That population was by no means pristine: it had been fished commercially for more than three centuries.

Our estimate of historic biomass revealed defining characteristics of past oceans. Nevertheless, it was so far from twenty-first century biomass estimates for the same region that it seemed stratospheric, somehow not to scale. Canadian managers today use a 1980s biomass estimate as their target for rebuilding cod. Many managers consider historical evidence as not "an appropriate reference point for present-day manag-




FIGURE 4. Above: Nova Scotian Shelf Catch Per Unit Effort 1852-1859. (CPUE = hundreds fish/day per vessel ton). Below: Estimates of cod biomass on the Nova Scotian Shelf over 167 years. **Adapted from Rosenberg, et. al., 2005, Frontiers in Ecology and the Environment.**

ers working to balance stock rebuilding with fisheries yield." Too many changes have occurred, they argue, which "preclude rebuilding a stock to its unfished level" (Henderschedt, 2015).

On the one hand, marine ecosystems are dynamic and susceptible to significant reconfiguration. Systems have changed. On the other hand, historical evidence provides perspective. The cod biomass figure for 1852 is a nagging reminder of how much coastal ecosystems have diminished in a very short time—only six human generations.

Our off-the-charts cod biomass estimate for 1852 illuminates the shifting baseline syndrome identified by Daniel Pauly in 1995. Pauly recognized that fishing pressure through time reduced ecosystem

complexity. Fishing shrank abundances, altered population structure by removing large individuals, affected predator-prey relationships, and changed fishes' geographic distribution. As a result, each human generation came to expect less, settling for a radically diminished natural world (Pauly, 1995).

It seems probable that cod's downturn during the 1850s was caused by synergy between overfishing and natural fluctuations. The middle of the nineteenth century was exceptionally cold, the final shudder of the Little Ice Age. Cold seawater can inhibit cod reproduction. As temperatures fell, North Atlantic cod productivity probably fell as well. Meanwhile, fishermen continued to hit stocks hard (Leavenworth, 2006; Rose, 2007).

As cod catches plummeted off-

shore during the 1850s, and as outraged inshore fishermen protested the new menhaden fishery that was destroying forage fish, animated discussions from Massachusetts to Nova Scotia focused on depletion and the need for conservation.

Using 1,664 inshore fishing logbooks from New England, along with an analysis of the entire fleet's tonnage and fishing patterns, our group was able to create a mathematically defensible landings estimate for Gulf of Maine cod in 1861. This predated all other data sets for that fishery. Our estimate, expressed as a range (depending on the multiplier used to convert whole fish to dried fish), was between 62,600 and 78,600 metric tons. Gulf of Maine landings have never been as good (Alexander et al., 2009).

Our landings estimate suggests that the fishery was thriving in 1861. Contemporaries did not think so. During the 1860s, when the average inshore boat was landing almost 16,000 cod per season, Maine fishermen proposed various bills to the legislature to reduce overfishing and save their cod. Conservation sentiment aimed at mackerel, menhaden, and cod was at its height among New England fishermen during the 1850s, 1860s, and 1870s. Yet despite those efforts, the situation worsened.

During the late nineteenth century unprecedented collapses occurred in four American fisheries - menhaden, mackerel, halibut, and lobster. They triggered protective legislation, bankrupt-

cies, and ecological havoc. Most striking is that only one of the four species, mackerel, had been fished commercially prior to the early nineteenth century. As Americans developed new markets and new technologies, previously underutilized species became desirable. Several were driven to the brink in a very short time.

Whale oil landings flattened out at mid-century because sailing ship technology had harvested most of the whales it could reach. "Whether Leviathan can long endure so wide a chase, and so remorseless a havoc," Herman Melville wrote in Moby-Dick in 1851, was the ques-




FIGURE 5A. Frenchman's Bay, Mt. Desert Island, Maine, 1885. U.S. Coast and Geodetic Survey.

tion, or "whether he must not at last be exterminated from the waters, and the last whale, like the last man, smoke his last pipe, and then himself evaporate in the final puff."

But oil could be rendered from menhaden as well as whales. America's first mechanized fishing boats were menhaden seiners, built in the 1870s. Menhaden landings soared, nearly all from inshore fisheries. Landings in 1878 surpassed those in many of the next sixty years, although catching technology got progressively more efficient, with larger ships, stronger nets, hydraulic haulers, and spotter aircraft.

Disaster struck in 1879. Menhaden barely appeared north of

their jobs. Good data exists on it reveals that menhaden populaand fishermen that the sea's bounty was not limitless.

Cape Cod that summer, and they were extraordinarily scarce for six years following. Oil factories closed. One thousand men lost menhaden landings from 1873 to the present. It is one of the longest time series in fisheries history, and tions fluctuate widely (Vaughan and Smith, 2009). Was the crash in 1879 a natural downturn, or the result of overfishing, or synergy between the two? We will never know for sure, though it set the stage for subsequent collapses, alerting politicians, industrialists,



THE 1880s mackerel was America's most popular food fish In 1884 mackerel

landings broke all records. This fishery was still conducted entirely under sail. Powerful modern mackerel schooners had great speed and windward ability.

Disaster struck in 1886. Fishermen landed less mackerel that year than in any of the previous fortyfive years. Bankruptcies followed. The mackerel crash resulted in the United States' first federal fishery law. In 1887 Congress closed the mackerel fishery for five years between March 1 and June 1, then understood as spawning season (Bolster, 2012). Mackerel landings would not reach their 1884 level again until 1968. By then, of course, motorized fishing ships were bigger, stronger, and more efficient.




FIGURE 5B. Scatterplot of aggregated cod catch per day in Frenchman's Bay in 1861. Almost all commercial fisheries were once conducted inshore. In Frenchman's Bay alone 148,704 cod were caught in 1861. Fisheries restoration policy should pay attention to spatial distribution as well as to overall numbers. Courtesy of Karen E. Alexander.

HE ATLANTIC halibut slaughter proceeded like that of American bison, ruthlessly and in plain sight. Halibut had been ignored for centuries, until the 1830s, when entrepreneurs began to promote them. During the 1840s

Gloucester created a halibut fleet.

In 1848, Atlantic halibut landings



FIGURE 6. Simple technologies affected fish stocks. Rowboats like these were central to American fishing prior to World War One, and they persisted longer in some fisheries. Upper: a seine boat used to encircle a school of mackerel or menhaden with a purse seine. Seine boats were towed behind schooners. Lower: a dory, which could be used to long-line cod, set lobster traps, or assist a seine boat crew. **Models by Erik Ronnberg. Photo courtesy of Erik Ronnberg.**

were approximately 20 million pounds. From the bonanza in the late 1840s, landings tailed off, and then fell precipitously. Shortly before 1900 they fell to about 9 million pounds; then to 3 million in 1910, and to about 1.25 million in 1915. Mopping up operations continued for a few more decades.

Today Atlantic halibut are so depleted from overfishing that they are off-limits to commercial fishing in American waters. It had taken only several human generations to destroy the population of a huge, well-known apex predator. During the 1880s Captain Joseph W. Collins lamented in a widely circulated U.S. Fish Commission publication that "if the present style of fishing is pursued" halibut "will in a few years become extremely scarce, if not almost extinct." He was right. And they were all hookcaught from small sailboats and rowboats (Bolster, 2012).

Lobsters were next. Everyone in the lobster business believed that the largest harvests occurred during the early 1870s. Maine's lobster landings were not tallied reliably until 1880. Nine years later saw the highpoint—24,451,219 pounds (Historical Maine Fisheries Landings Data).

Disaster struck in the 1890s. Landings fell precipitously that decade, and then fell again. Not until 1957 would Maine lobstermen land as much as had been landed 68 years earlier. By then lobstermen had large motorized boats, with mechanical haulers, and nearly five times as many traps to catch the same weight of lobsters that had been caught in 1889, when men fished shallow waters from sailboats and rowboats. Fool-hardy overfishing at mid-century decimated the lobster fishery (http://www.maine.gov/ dmr/commercial-fishing/landings/

documents/lobster.table.pdf).

Unlike halibut, however, lobster populations rebounded toward the end of the twentieth century, and in the new millennium they have been astronomical. 2016 saw a record harvest by Maine lobstermen: approximately 131 million pounds. Today Gulf of Maine fishermen have nearly all their eggs in one basket—the lobster fishery. About 80% of Maine's fishing revenue now is lobster based. When the stock crashes it will create a catastrophe far worse than that of the 1890s (http://www.maine.gov/ dmr/commercial-fishing/landings/ documents/lobster.table.pdf).

Several points from this brief narrative deserve attention. All of those fisheries were initially inshore operations (Figure 5). As depletion ensued, fishermen moved offshore. Recent management plans have paid insufficient attention to fishes' spatial distribution. But it matters. Another point, too often ignored, is that prior to industrialized fishing, humans wielding very simple technologies affected marine ecosystems in profound ways (Figure 6). Finally: fluctuations in those systems were the norm. Human impacts must be assessed against constantly occurring natural changes.

Ill-advised as were the post-Civil War halibut and lobster industries, reckless abandon was not the norm in every fishery. What is most striking about those years was fishermen's insistence that the resources on which they depended were shrinking, and their realization that new gear and new fisheries were exacerbating the problem.

In 1861 a legislative committee in Nova Scotia came down hard on long-lining, a new technology with considerably more catching power than old hand-lines. It is well known, they wrote, that "if this mode of taking fish" continues, "in a few years these banks ... will be rendered altogether unproductive." The next year legislators in both Newfoundland and Maine introduced bills to prohibit long-lining. Throughout the 1860s numerous attempts to prevent long-lining "for the purpose of protecting the cod fishery" were introduced in coastal legislatures. In 1869 Massachusetts' fish commissioners noted complaints "of a diminution in many species" (Bolster, 2012).

Here is the tragedy: Fishermen often vehemently protested new gear, convinced it would drive a nail into the coffin of the fisheries—though they ultimately adopted it. That's the shifting baseline

syndrome. Time and time again, knowledge of previous depletion was lost with adoption of more efficient gear. In 1870, for instance, the Portsmouth, New Hampshire cod fishery was thriving: according to a newspaper "over a million pounds of codfish have been landed at one wharf in Portsmouth during the past winter." Despite their concerns, local fishermen had adopted long-lines. "In and about the harbor, there is now sunk over 63 miles of trawls [long-lines], on which are hung over 96,000 hooks." Not long before, such gear would have been considered immoral. By 1870 it was the new norm (Gloucester Telegraph, 1870).

1905 saw introduction of the first steam-powered otter trawler (bottom dragger) in the western Atlantic, the Spray (Figure 7). A close copy of British fishing ships that had redefined North Sea fishing, Spray actively pursued fish by towing a net on the bottom. That was revolutionary. Fishermen had always waited for fish to come to hooks or gill-nets (Figure 8).

"The time to stop this thing is while it is in its beginning," insisted a Massachusetts Congressman, who introduced a bill in 1911 to prohibit trawling. The Gloucester Board of Trade backed the prohibition, citing evidence of North Sea depletion by fleets of steam trawlers. John F. Fitzgerald, the mayor of what he called "the biggest fish port in the western world" (Boston), expressed concerns about "wiping out the fish industry." Colossal opposition to



FIGURE 7. Most fishermen initially feared destructive trawling technology and resisted it. The first encounter of schooner fishermen with a steam trawler in the western hemisphere took place shortly after the launch of Spray, in 1905. Thomas M. Hoyne, New Ways on Quero Bank, 1981; courtesy of Doris O. Hoyne and the Peabody Essex Museum, Salem, Mass. Gift of Russell W. Knight, 1982.

bottom trawling existed, based on insiders' knowledge that fisheries were already depleted. But Congress refused to ban the new technology (Bolster, 2012).

By 1914 the U.S. Fish Commission (and its successor, the Bureau of Fisheries) had existed for more than forty years, spending considerable taxpayer dollars to increase landings. Among other tactics, they managed an aggressive program for propagating fish, clams, and lobsters. Scientists liked to laud their accomplishments, but remaining optimistic was not easy, even before a New York Times reporter drew on decades of Bureau of Fisheries data for a major feature in 1914.

"Extermination Threatens American Sea Fisheries - Cost to Consumer Has Risen between 10 and 600 Per Cent Because of Decrease in Supply" (Widenmann, 1914) (Figure 9). The Bureau's storied efforts to propagate sea fish had not worked. The defining trajectory was clear, and disconcerting. Fishermen went farther afield, fished at greater depths, but brought home less. Draggers were just coming on line. An avalanche of cheap fish would soon silence the critics, camouflaging the extent of damage already done.

Most people today know at least the rough outlines of what happened next. Bottom trawling initially caught lots of fish, including unwanted ones. By-catch soared, as did habitat destruction Fishing boats got bigger. Technology to find and catch fish became more sophisticated. Western Atlantic catches ultimately peaked during the late 1960s, with huge foreign factory ships. By 1976, American and Canadian ground fish had been decimated. A further

low point came in the 1990s. Slight rebuilding followed. We are living through another low point now.

The latest tragedy is that managers' efforts to reduce fishing pressure and rebuild Gulf of Maine




FIGURE 8. Otter trawls were revolutionary. Fishermen no longer waited for fish to approach a hook or gill net, but actively pursued them. In spite of serious opposition, by about 1925 otter trawling was the new "normal." Fishermen who had long feared its destructiveness came to embrace it, becoming victims of the shifting baseline syndrome. From John Dyson. Business in Great Waters: The Story of British Fishermen (London, 1977), 261.




FIGURE 9. Based on forty years of data collected by the U.S. Fish Commission and the Bureau of Fisheries, this article revealed that while catching power had expanded, catches continued to decline. Revolutionary new bottom trawling technology would soon obscure that chilling insight with an avalanche of cheap fish. From The New York Times, July 26, 1914.

cod have been undermined, apparently by warming water. Warming water has reduced recruitment and increased mortality in cod stocks (Pershing et al., 2015). Once again, marine systems' complexity and dynamism has thwarted the best efforts to manage those systems for humans' desired outcomes.

Swordfish have been one of management's great successes in the last quarter century, though historical evidence suggests that swords' recovery has not been as extensive as many believe. From 1960 to 1996, stocks declined 68 percent, while the average size of North Atlantic swordfish dropped from 266 to 90 pounds. During the 1990s the U.S. reduced landings to rebuild stocks. The International Commission for the Conservation of Atlantic Tuna (ICCAT)—the regional fisheries management organization tasked with managing high seas swordfish—adopted a rebuilding plan. Swordfish grow quickly, and spawn in temporallyand spatially-dispersed ways. With less fishing pressure, swordfish rebounded. An assessment in 2009 found the stock rebuilt. An assessment in 2013 concluded that its biomass would allow sustainable fishing (Neilson et al., 2013; http:// www.nmfs.noaa.gov/ia/slider_stories/2013/11/swordfish.html).

Considerable historical evidence from 80 to 130 years ago, however, structured into quantitative analyses, shows that swordfish were then abundant in nearshore regions of New England and Nova




FIGURE 10. Historical approaches provide sorely needed perspective. This graph is not precise, but it is accurate. Courtesy Karen E. Alexander

swordfish biomass is now at a sustainable level, swords have not repopulated their historic range. Moreover, captured fish today are much smaller. Technological shifts in the fishery have allowed fishermen to pursue them hundreds of miles offshore, and landings remain robust. But the current landings data and biomass estimates about which regulators are so happy exist over a huge spatial scale the stock's entire range. Closegrained historical data reveal local depletions in areas where swordfish were once common. Shifted spatial baselines may be obscuring the extent of the stock's rebuilding (MacKenzie et al, in preparation).

One waypoint remains relatively known in the rancorous discussions that have characterized American fisheries policy for the last 160 years. From the 1850s through the 1920s, fishermen la-

Scotia. Although North Atlantic mented depletions. Using terms such as "diminution" and "degradation," they begged politicians to solve the fisheries problem. Scientists, meanwhile, led by the U.S. Fish Commission, generally encouraged more robust fishing, believing that the biological knowledge and better fishing gear they were developing would sustain larger harvests.

> During the early 1930s, Woods Hole biologists revealed notoriously poor year classes of haddock on Georges Bank. The scientific community recommended larger mesh size and smaller harvests. That was radical—one of the first instances in which fisheries scientists suggested reducing catches. By then bottom trawling had become the new norm, and fishermen—having lost sight of previous depletions had thrown caution to the wind. American fishermen reacted angrily to the idea of throttling back,

convinced that if they did not catch those haddock, Canadians would.

The terms of the debate remained the same: "Not enough fish" versus "fish harder." But the sides switched. Fishermen, who had long lamented depletion, insisted during the 1930s they should fish without restriction. Scientists, who had long believed in the sea's ultimate productivity, began to advocate conservation. Both sides continued to talk past each other (Bolster, 2012).

Discussions of the problem have persisted for more than 160 years, even as the problem has worsened. Is it time to take a different tack? Or should we just count the few fish that remain more carefully (Figure 10)?



Looking Forward

OMPLEX systems deserve complex analysis. Without discarding statistical stock analysis, despite its limitations and uncertainties, we might incorporate other forms of knowledge into management plans, including the contextualization and perspective that history provides.

We should develop techniques to incorporate historical information, when possible, as an additional data source for use in reconstructing aspects of past ecosys-

tems. For instance, the presence, absence, and descriptions of organisms in particular times and at particular locations have long been grouped, ranked, ordered, and analyzed to show decline. Images of species' size and abundance have likewise been analyzed (Jackson, 1997; McClenachan and Cooper, 2008; McClenachan, 2009). Historical evidence can be binned at intervals. Terms used repetitively can reveal relative abundance. We found 3028 descriptions, in seven categories, ranking cod abundance in nineteenth-century Frenchman's Bay logbooks: no fish, fish very scarce, fish scarce, fish, some fish, fish plenty, and fish very plenty. Correlations existed between cod rank and cod landings. Similar descriptions of bait abundance provided more ecosystem indicators. Arithmetic, simple statistics, and GIS applied to historical observations hold out the promise of creating snapshots from the past (Alexander et al., 2009).

Binning evidence historically has shown how humans have influenced marine ecosystems over long periods of time (Pandolfi, 2003; Lotze and Milewski, 2004), and that an extreme climate event in 1815 permanently transformed Gulf of Maine fisheries (Alexander et al., 2017).

Even if historical evidence is not fully integrated into scientific data sets, it could frame data and results. According to NOAA's Fish Watch, for instance, haddock is fully recovered, and not

overfished. Moreover, overfishing is not occurring. In light of depleted haddock stocks in the early 1990s, the recent spawning stock biomass of Gulf of Maine haddock reflects managers' success - 10,325 mt in 2015. Haddock has rebounded dramatically. Historical data, however, provide perspective. In 1902, almost 10,000 mt of haddock were caught in the Gulf of Maine. That makes me question the assertion that haddock is "fully recovered" (https:// www.nefsc.noaa.gov/groundfish/ operational-assessments-2015/ Reports/2015 HAD GM Assessment_Report_20150913.pdf; Alexander et al., 2011).

A NOAA report for all managed fisheries stated "39 Stocks Rebuilt as of December 31, 2015." Many pronouncements about "rebuilt" stocks are based on severely limited management targets, not on historical ecology (http://www.nmfs.noaa. gov/sfa/fisheries_eco/status_of_fisheries/archive/2015/fourth/rebuilt stocks_q4_2015.png).



How might management change?

O BEGIN with, policy could catch up to the best available science. Considerable research in marine historical ecology and marine environmental history has worked its way into the literature in the last twenty years. Policy-makers committed to sustainability and resilience could draw on that work. Ultimately, of course, it will not be honored sufficiently until standards about "the best scientific information available" are revised to include observations from the past (Engelhard et al., 2016).

Conservation advocates have suggested more Marine Protected Areas, seasonal moratoriums, and promoting recovery of forage fish. Historians and historical ecologists concur, but suggest as well that we learn to manage with an eye on changes over time, as well as an eye on spatial distribution within fishes' range. The entire range of a species is not a meaningful unit of analysis. Let's pay attention to where fish were caught, and where they are caught. Can we create policy that promotes not just recovery of biomass, but restoration of fishes' historic ranges (Alexander et al., 2011)?

Given the devastated state of marine ecosystems, perhaps we should be managing for resilience, rather than for maximum sustainable yield. The stated purpose of the Magnuson-Stevens Act is to "promote domestic and commercial recreational fishing under sound conservation and management principles" through "the best scientific information available." I would like to see the law flipped to emphasize conservation and ecosystem recovery, while allowing some fishing. I know that is an unrealistic aspiration at this time, but so was the abolition of slavery during the 1840s. We must live in hopes.

Given that marine ecosystems are coupled human-natural systems, we should assess them holistically. That is clearly the goal of ecosystem-based management (Pikitch, et al., 2004). I'm encouraged, but leery. Ecosystem based management may require more data to build even more complicated models. If current standards regarding data acceptability remain in place, historical evidence will continue to be excluded.

silos remain a problem. I would like to see rising generations of scientists and scholars trained in interdisciplinary ways, so that perceptions are broadened.

Finally, we must give this process time. Rebuilding will not occur in the two-year or fouryear political cycle, or in the ten years stipulated by the Magnuson-Stevens Act. The National Marine Fisheries Service is charged with managing the tattered remnant of a once-productive system. Given the system's fragility, and on-going environmental change, further collapses are likely (Pershing et al., 2015). Recovery may be possible, or it may not. To find out, we need to hunker down for a long haul with a precautionary approach.

Fisheries history reveals several bold lessons. First: humans have limited ability to control nature. Second: Humans have an es-

tablished record of fouling up the ecosystems on which we rely for crucial goods and services. That sort of framing deserves attention. It would help us link management with humility. The bottom line is that we take too much. That has been the case for centuries, and well known, too. It's not just fish Look at water in the American West (Reisner, 1986).

My immersion in historical fisheries was sobering. Yet my intimacy with depictions of nearly unfathomable abundance also promoted a sense of wonder. That inspiration is worth nurturing, and Self-perpetuating academic right now it is only available in the historical record.



HANKS TO Molly Bolster, Karen E. Alexander, Emily Klein, Matthew G. McKenzie, Susan Roberts, Stacee Karras, and Steve Jones for thoughtful reading of this manuscript. My training as a marine environmental historian grew out of the Census of Marine Life, specifically its History of Marine Animals Project (HMAP), and the UNH Cod Project. We received substantial funding from the Alfred P. Sloan Foundation, NSF, NH Sea Grant, NOAA's Marine Sanctuary Program, and a UNH Center for

the Humanities Senior Fellowship.

SOURCES CITED | Al-Abdulrazzak, A., Zeller, D., and Pauly, D., 2015. Understanding fisheries through historical reconstructions. Marine Historical Ecology in Conservation, eds. Kittinger, J.N., McClenachan, L., Gedan, K., and Blight, L.K.. | Alexander, K.E., Leavenworth, W.B., Cournane, J., Cooper, A.B., Claesson, S., Brennan, S., Smith, G., Rains, L., Magness, K., Dunn, R., Law, T., Gee, R., Bolster, W.J., Rosenberg, A.A., 2009. Gulf of Maine Cod in 1861: historical analysis of fishery logbooks, with ecosystem implications. Fish and Fisheries 10, 428-449. | Alexander, K.E., W.B. Leavenworth, S. Claesson, W.J. Bolster. (2011) Catch Density: a new approach to shifting baselines, stock assessment, and ecosystem-based management. Bulletin of Marine Science 87(2):213-234. | Alexander, K.E., Leavenworth, W.B., Willis, T.V., Hall, C., Mattocks, S., Bittner, S., Klein, E., 2017. Tambora and the mackerel year: phenology and fisheries during an extreme climate event. Science advances 3, no. 1, e1601635. | Appleby, J., Hunt, L., Jacob, M., 1994. Telling the Truth about History. | Benedict, R., 1943. Obituary, Franz Boas. Science 97, 60. | Bolster, W.J., 2006. Opportunities in Marine Environmental History. Environmental History 11, 567-597. | Bolster, W.J., Alexander, K.E., Leavenworth, W.B., 2011. The historic abundance of cod on the Nova Scotian Shelf in Shifting Baselines: The Past and Future of Ocean Resources, eds. Jackson, J.B.C., Alexander, K.E., Sala, E. | Bolster, W. J., 2012. The Mortal Sea: Fishing the Atlantic in the Age of Sail. | Canada Department of Fisheries and Oceans, 2002. Science Stock Status Report for Cod Areas 4X and 4VsW. www.mar.dfo-mpo.gc.ca/science/rap/internet/. | Canada Department of Fisheries and Oceans, 2017. 2016 4X5Yb Atlantic Cod (Gadus morhua) Stock Status Update. DFO Can. Sci. Advis. Sec. Sci. Resp. 2017/024. | Chadbourne, B., c. 1797. "A Description of the Town and Village." Mss., Old Berwick Historical Society, South Berwick, Maine. | Engelhard, G., Thurstan, R.H., MacKenzie, B.R., Alleway, H.K., Colin, R., Bannister, A., Cardinale, M., Clarke, M.W., Currie, J.C., Fortibuoni, T., Holm, P., Holt, S.J., Mazzoldi, C., Pinnegar, J.K., Raicevich, S., Volckaert, F.A.M., Klein, E., Lescrauwaet, A-K., 2016. ICES meets historical marine ecology: placing the history of fish and fisheries in current policy context. ICES Journal of Marine Science, doi: 10.1093/icesjms/fsv219. | Franklin, H.B., 2007. The most important fish in the sea. | Gloucester Telegraph, March 23, 1870. | Henderschedt, J., 2015. Viewpoint from a practitioner: Real-world constraints to applying historical baselines in U.S. fisheries. Marine Historical Ecology in Conservation. | Historical Maine Fisheries Landings Data, http://www.maine.gov/ dmr/commercial-fishing/landings/historical-data.html | Jackson, J.B.C., Reefs Since Columbus. Coral Reefs 16 (Supplment), S23-S32. | Jackson, J.B.C., Kirby, M.X., Berger, W.H., Bjorndal, K.A., Botsford, L.W., Bourque, B.J., Bradbury, R.H., Cooke, R., Erlandson, J., Estes, J.A., Hughes, T.P., Kidwell, S., Lange, C.B., Lenihan, H.S., Pandolfi, J.M., Peterson, C.H., Steneck, R.S., Tegner, M.J., Warner, R.R., 2001. Historical overfishing and the recent collapse of coastal ecosystems. Science, 293, 629-638. Kittinger, J.N., McClenachan, L., Gedan, K.B., Blight, L.K., 2014. Marine Historical Ecology in Conservation: Applying the Past to Manage for the Future. Oakland, CA: University of California Press. | Leavenworth, W.B., 2006. Opening pandora's box: tradition, competition, and technology on the Scotian Shelf, 1852-1860, in The North Atlantic Fisheries: Supply, Marketing, and Consumption, 1560-1990. Eds. Starkey, D.J. and Candow, J.E. I Link, J.S., Ihde, T.F., Harvey, C.J., Gaichas, S.K., Field, J.C., Brodziak, J.K.T., Townsend, H.M., Perman, R.M., 2012. Dealing with uncertainty in ecosystem models: the paradox of use for living resource management. Progress in Oceanography 102, 102-114. | Liu J., Dietz, T., Carpenter, S.R., Alberti, M., Folke, C., Moran, E., Pell, A. N., Deadman, P., Kratz, T., Lubchenco, J., Ostrom, E., Ouyang, Z., Provencher, W., Redman, C.L., Schneider, S. H., Taylor, W.W., 2007. Complexity of Coupled Human and Natural Systems. Science, 317, 1513-1516. | Lotze, H.K., Milewski, I., 2004. Two centuries of multiple human impacts and successive changes in a north Atlantic food web. Ecological Applications 14, 1428-1447. | Lotze, H.K., Lenihan, H.S., Bourque, B.J., Bradbury, R.H., Cooke, R.G., Kay, M.C., Kidwell, S.M., Kirby, M.X., Peterson, C.H., Jackson, J.B.C., 2006. Depletion, Degradation, and Recovery Potential of Estuaries and Coastal Seas. Science 312,1806-1809. | Lowenthal. D., 1985. The Past is a Foreign Country. | MacKenzie, B., Alexander, K.E., Leavenworth, W.B., Claesson, S., Bolster, W.J., and Cooper, A., (in preparation). Historical evidence opens new swordfish recovery perspectives in the northwest Atlantic. | McClenachan, L., Cooper, A.B., 2008. Extinction rates, historical population structure and ecological role of the Caribbean monk seal. Proceedings of the Royal Society B, 275, 1351-1358. | McClenacan, L., 2009. Documenting loss of large trophy fish from the Florida Keys with historical photographs. Conservation Biology, 23, 636-643. | McClenachan, L., Lovell, S., Keaveney, C., 2015. Social benefits of restoring historical ecosystems and fisheries: alewives in Maine. Ecology and Society 20 (2): 31. | MeSA, Maine State Archive, Legislative Graveyard, 1857. Petition of inhabitants of Gouldsboro, Oct. 10, 1857, box 256, folder37. | MeSA, Maine State Archive, Legislative Laws, 1852. Petition of inhabitants of Southport, Boothbay, and vicinity, Jan., 1852, box 265, folder 125. | Meyers, R., and Worm, B., 2003. Rapid world-wide depletion of predatory fish communities. Nature 423, 280-283. | National Research Council (NRC), 2004. Improving the Use of the "Best Scientific Information Available" Standard in Fisheries Management. Washington, DC: National Academies Press. | Neilson, J., Arocha, F., Cass-Calay, S., Mejuto, J., Ortiz, M., Scott, G., Smith, C., Travassos, P., Tserpes, G., and Andrushchenko, I., 2013. The Recovery of Atlantic Swordfish: The Comparative Roles of the Regional Fisheries Management Organization and Species Biology. Reviews in Fisheries Science, 21 (2), 59-97. | Pandolfi, J.M., Bradbury, R.H., Sala, E. et al. 2003. Global trajectories of the long-term decline of coral reef ecosystems. Science, 301, 955-958. | Pauly, D., 1995. Anecdotes and the shifting baseline syndrome of fisheries. Trends in Ecology and Evolution, 10, 430. | Pauly, D., and Maclean, J., 2003. In a Perfect Ocean: The State of Fisheries and Ecosystems in the North Atlantic Ocean. | Pauly, P.J., 2000. Biologists and the Promise of American Life: From Meriwether Lewis to Alfred Kinsey. | Pershing, A.J., Alexander, M.A., Hernandez, C.M., Kerr, L.A., LeBris, A., Mills, K.E., Nye, J.A., Record, N.R., Scannell, H.A., Scott, J.D., Sherwood, G.D., Thomas, A.C., 2015. Slow adaptation in the face of rapid warming leads to collapse of Gulf of Maine cod industry. Science. 2015 Nov 13;350(6262):809-12. doi: 10.1126/science.aac9819. | Pikitch, E.A., Santora, C., Babcock, E.A., Bakun, A., Bonfil, R., Conover, D.O., Dayton, P., Doukakis, P., Fluharty, D., Heneman, B., Houde, E.D., Link, J., Livingston, P.A., Mangel, M., McAllister, M.K., Pope, J., and Sainsbury, K.J., 2004, Ecosystem-Based Fishery Management, Science, 305, 346-347, I Reisner, Marc. 1986, Cadillac Desert: The American West and its Disappearing Water. | Rose, G.A., 2007. Cod: The Ecological History of the North Atlantic Fisheries. | Rosenberg, A.A., Bolster, W. J., Alexander, K.E., Leavenworth, W.B, Cooper, A.B., McKenzie, M.G., 2005. The History of Ocean Resources: Modeling Cod Biomass Using Historical Records. Frontiers in Ecology and the Environment, 3, 84-90. | USCFF, 1874. Report of the Commissioner for 1872 and 1873, xiv. | Vaughan, D.S. and Smith, J.S., 2009. Reconstructing Historical Commercial Landings of Atlantic Menhaden. SEDAR 20-DW02. | Warner, W., 1977. Distant Water: The Fate of the North Atlantic Fisherman, I Widenmann, R.A., 1914. Extermination Threatens American Sea Fishes. New York Times, July 26, 1914.



