

International
Muon Collider
Collaboration



Muon Collider Collaboration

D. Schulte
for the International Muon Collider Collaboration

EPP2024
December 2022

Motivation and Goal



Previous studies in US (now very strong interest again), experimental programme in UK and alternatives studies by INFN

New strong interest in **high-energy, high-luminosity lepton collider**

- Combines **precision physics** and **discovery reach**

Muon collider promises **sustainable** approach to the **energy frontier**

- limited power consumption, cost and land use

Technology and **design advances** in past years

- review did not find any showstoppers

Goal is


- 10+ TeV collider
- potential initial energy stage (e.g. 3 TeV)
- higher energies to be explored later

$\sqrt{s}$	$\int \mathcal{L} dt$
3 TeV	1 ab ⁻¹
10 TeV	10 ab ⁻¹
14 TeV	20 ab ⁻¹

A new Interest in Muon Colliders

From, e.g., Snowmass21 EF report draft:

"A 10-TeV scale muon collider with sufficient integrated luminosity provides an energy reach similar to that of a 100 TeV proton-proton collider. [...] muon and hadron colliders have similar reach and can significantly constrain scenarios motivated by the naturalness principle. [...] Multi-TeV muon colliders will have the benefit of excellent signal to background [...] One of the key measurements from the multi-TeV colliders is the one of the Higgs self-coupling to a precision of a few percent, and the scanning of the Higgs potential."



Fabio Maltoni - Physics



from F. Maltoni at IMCC Annual Meeting




A. Wulzer, F. Maltoni, P. Meade et al.

O(150) authors, 15 editors, 100 papers



Selected summary plots, from Snowmass21 reports:

2 IMCC reports, plus Muon Collider Forum report. Total of 15 editors, ~150 authors, based on ~100 papers from 3 past years



Strong US involvement starting with **Muon Smasher's Guide** and in **Muon Collider Forum**

Cost and Sustainability



CLIC is highest energy proposal with CDR

- No obvious way to further improve linear colliders (decades of R&D)
- Cost 18 GCHF, power approx. 500 MW

Rough rule of thumb:

- cost proportional to energy
- power proportional to luminosity

Muon Collider goals (10 TeV), challenging but reasonable:

- Much **more luminosity** than CLIC at 3 TeV ($L=20 \times 10^{34}$, CLIC: $L=2 \times 10^{34} / 6 \times 10^{34}$)
- **Lower power consumption** than CLIC at 3 TeV ($P_{beam,MC}=0.5 P_{beam,CLIC}$)
- **Lower cost**


Staging is possible

Synergies exist (neutrino/higgs)

Unique opportunity for a **high-energy, high-luminosity lepton collider**

Technically Limited Timeline

To be reviewed considering progress, funding and decisions



Muon collider important in the long term

Prudently explore if MuC can be **option as next project**

- e.g. in Europe if higgs factory built elsewhere
- **sufficient funding required now**
- very **strong ramp-up required** after 2026
- might require compromises on initial scope and performance
 - e.g. 3 TeV

Accelerator R&D Roadmap



No showstopper identified by Roadmap or Implementation Task Force

- mainly technological challenges
- details in Roadmap

Full funding scenario deliverables by next ESPPU/other processes

- **Project Evaluation Report**
 - key performance, risk, cost and power drivers
 - site considerations (CERN and elsewhere)
- **R&D Plan**
 - describes a path towards the collider;
 - key element is **demonstrator concept**
- **Interim Report (2023)**

Allows to make **informed decisions**

Current funding level allows only to address the most critical items

- making priorities based on risk and collaborator interest

<http://arxiv.org/abs/2201.07895>

Label	Begin	End	Description	Aspirational		Minimal	
				[FTEy]	[kCHF]	[FTEy]	[kCHF]
MC.SITE	2021	2025	Site and layout	15.5	300	13.5	300
MC.NF	2022	2026	Neutrino flux mitigation system	22.5	250	0	0
MC.MDI	2021	2025	Machining-detector interface	15	0	15	0
MC.ACC.CR	2022	2025	Collider ring	10	0	10	0
MC.ACC.HE	2022	2025	High-energy complex	11	0	7.5	0
MC.ACC.MC	2021	2025	Muon cooling systems	47	0	22	0
MC.ACC.P	2022	2026	Proton complex	26	0	3.5	0
MC.ACC.COLL	2022	2025	Collective effects across complex	18.2	0	18.2	0
MC.ACC.ALT	2022	2025	High-energy alternatives	11.7	0	0	0
MC.HFM.HE	2022	2025	High-field magnets	6.5	0	6.5	0
MC.HFM.SOL	2022	2026	High-field solenoids	76	2700	29	0
MC.FR	2021	2026	Fast-ramping magnet system	27.5	1020	22.5	520
MC.RF.HE	2021	2026	High Energy complex RF	10.6	0	7.6	0
MC.REMC	2022	2026	Muon cooling RF	13.6	0	7	0
MC.RF.TS	2024	2026	RF test stand + test cavities	10	3300	0	0
MC.MOD	2022	2026	Muon cooling test module	17.7	400	4.9	100
MC.DEM	2022	2026	Cooling demonstrator design	34.1	1250	3.8	250
MC.TAR	2022	2026	Target system	60	1405	9	25
MC.INT	2022	2026	Coordination and integration	13	1250	13	1250
Sum				445.9	11875	193	2445

Table 5.5: The resource requirements for the two scenarios. The personnel estimate is given in full-time equivalent years and the material in kCHF. It should be noted that the personnel contains a significant number of PhD students. Material budgets do not include budget for travel, personal IT equipment and similar costs. Colours are included for comparison with the resource profile Fig. 5.7.

Collaboration Vision




IMCC is an **international** collaboration and aims to

- Enlarge the collaboration
 - Physics interest in all regions, strong US contribution to the muon collider physics and detector, interest in Japan
 - First US university have joined collaboration, try to see how to move forward, also with labs
- Combine the R&D efforts for the design and its technologies
 - Critical contributions in all relevant fields by the US
- Consider several sites for the collider
 - CERN would be one, FNAL and others should also be considered
 - A proposal with alternative sites is stronger for a single site
- Consider several sites for the demonstrators
 - E.g. Muon production and cooling demonstrator at CERN, FNAL, ESS, JPARC
 - e.g. RCS at ESRF or elsewhere
 - Target tests
 - ...

Collaboration Organisation

- **Collaboration Board (ICB)**
 - Chair: **Nadia Pastrone**
- **Steering Board (ISB)**
 - Chair **Steinar Stapnes**
 - Reports to LDG but could add DOE
- **Advisory Committee (IAC)**
 - To be defined
- **Coordination committee (CC)**
 - Study Leader **Daniel Schulte**
 - Deputies: **Andrea Wulzer, Donatella Lucchesi, Chris Rogers**
 - **Sergo Jindariani, Mark Palmer** as US links
 - Will strengthen physics and detectors




Key Challenges

0) Physics case

4) Drives the **beam quality**
MAP put much effort in design
optimise as much as possible

2) Beam-induced
background



3) **Cost** and **power** consumption limit energy reach
e.g. 35 km accelerator for 10 TeV, 10 km collider ring
Also impacts **beam quality**

1) **Dense neutrino flux**
mitigated by mover system
and site selection

Muon Decay and Detector Background

Muons decay produces electrons and positrons

- Loss per unit length almost independent of energy

Tools mostly ready to generate background

- tentative beamline and mask, FLUKA
- tentative beam-beam for muons (GUINEA-PIG)


Studies at 1.5 and 3 TeV with concept based on CLIC detector

- **Radiation level in tracking detector similar to HL-LHC**


Studies with **beam-induced background** in progress

- some channels are not affected by background
- some improvement required for other channels

Concept for **10 TeV** in progress




Detector team
O(69) authors, O(150
signatories)




Neutrino Flux

Goal: **similar to LHC**: limit neutrino flux to have **negligible impact**, “fully optimised” (10% of MAP goal)
Verify performance of concept to be good for 14 TeV




FLUKA dose studies




G. Lerner, D. Calzolari,
A. Lechner, C. Ahdida

Conformity Verification Scheme




C. Ahdida, P. Vojtyla, M. Witorski, H. Vincke

Mover and support system




F. Bertinelli et al. (CERN, Riga)

G. Lacerda, Y. Robert, N. Guilhaudin (CERN)



Flux direction map / lattice design / mover impact on beam



C. Carli, K. Skoufaris (CERN)


Mitigation: Site choice tool

D. Schulte

Muon Collider, EPP2024, December 2022

High-energy Muon Complex

Acceleration Complex



Core of baseline is sequence of pulsed synchrotron (0.4-11 ms)
Important cost and power consumption

Started

- **Integrated design of RCS**
 - lattice with realistic hardware specifications
 - collective effects
- **Concept of key components**
 - Fast-ramping normal magnets
 - HTS alternative
 - Efficient power converters
 - RF with transient beam loading

A. Chance et al. (CEA)

E. Metral et al. (CERN)

L. Bottura et al. (LNCMI, Darmstadt, Bologna, Twente)


F. Boattini et al.

H. Damerell, F. Batsch, U. van Rienen, A. Grudiev et al. (Rostock, Milano, CERN)

- Alternative FFA S. Machida et al. (RAL)



FNAL 300 T/s HTS magnet



EMMA proof of FFAG principle
Nature Physics 8
23-247 (2012)

D. Schulte

Muon Collider, EPP2024, December 2022



Collider Ring

MAP developed 4.5 km ring for 3 TeV with Nb₃Sn
• magnet specifications in the HL-LHC range


Work progressing on 10 km ring for 10 TeV collider ring

- around 16 T Nb₃Sn or HTS dipoles
- final focus based on HTS

15 cm aperture for shielding to ensure magnet lifetime


Need stress managed magnet designs

INFN, Milano, Kyoto, CERN, profit from US



D. Schulte

Muon Collider, EPP2024, December 2022



C. Carli, K. Skoufaris (CERN)

Field choice will be reviewed for cost

Example alternatives:

- a 6 km 3 TeV ring with NbTi at 8 T in arcs
- a 15 km 10 TeV ring with HL-LHC performances
- slight reduction in luminosity

Hybrid RCS concept

- lattice, beam dynamics, magnets, power converter, superconducting RF

Collider ring concept

3 TeV MAP design exists

10 TeV more demanding, ongoing


D. Schulte

Muon Collider, EPP2024, December 2022

Muon Production and Cooling

Target

MAP target design, K. McDonald, et al.



2 MW proton beam is OK

Bunching challenge will be addressed by ESS experts

N. Milas et al. (ESS, Uppsala)


D. Schulte

Large bore, high field solenoid

Nb₃Sn or HTS

Similar to ITER


A. Lechner et al.
L. Bottura et al.



ITER Central Solenoid Model Coil
13 T in 1.7 m (LTS)

Shock and power in target:
Simulations of graphite target indicate 2 MW could be acceptable


STFC will also study alternatives



Muon Collider, EPP2024, December 2022



Muon Cooling



C. Rogers, B. Stechauner, E. Fol et al. (RAL, CERN)




MICE Collaboration

Nature vol. 578, p. 53-59 (2020)

Principle demonstrated with no RF
Use of data for benchmarking is still ongoing

MAP designs almost achieve 10 TeV goal
• miss factor two for final cooling



Integration/optimisation of design
Integrating improved technologies
Collective effects

C. Rogers et al. (RAL, CERN)

T. Pieloni et al. (EPFL, CERN)

Cooling Cell Technology

C. Marchand, Alexej Grudiev et al. (CEA, Milano, CERN, Tartu)


RF cavities in magnetic field

MAP demonstrated higher than goal gradient
Improve design based on theoretical understanding
Preparation of **new test stand**, but needs funding

- Test stand at CEA (700 MHz, need funding)
- Test at other frequencies in the UK considered
- Use of CLIC breakdown experiment considered

Assessment of realistic goal for highest field solenoids

- MAP demonstrated 30 T
- now magnets aim for 40+ T
- even more can be possible




L. Bottura et al. INFN (Task Leader), CEA, CERN, LNCMI, PSI, SOTON, UNIGE and TWENTE, in collaboration with KEK and US-MDP

Will develop example **cooling cell integration**

- tight constraints
- additional technologies (absorbers, instrumentation,...)
- early preparation of **demonstrator facility**

L. Rossi et al. (INFN, Milano, STFC, CERN), J. Ferreira Somoza et al.

D. Schulte



Muon Collider, EPP2024, December 2022

Limited efforts on

- proton complex
- target
- cooling channel optimisation
- cooling channel technologies

New is integrated cooling cell design

D. Schulte


Muon Collider, EPP2024, December 2022



International Muon Collider Collaboration



International Muon Collider Collaboration



MuCool demonstrated >50 MV/m in 5 T

- W-filled copper
- Be-end caps

Key Technologies



Magnets

- Superconducting solenoids for target and cooling profit from developments for society
 - target solenoid comparable to ITER central solenoid fusion
 - 6D cooling solenoids similar and wind power generators, motors
 - final cooling solenoids synergetic with high-field research, NMR
- Collider ring magnets
 - profit from developments for other colliders FCC-hh, stress-managed magnets
- Fast-ramping normal-conducting magnet system
 - HTS alternative, power converter

RF systems

- superconducting RF, normal-conducting RF, efficient klystrons

Target, cooling absorbers, windows, shielding



Neutrino mitigation mover system, cooling cell integration, ...

Detector

CDR Phase

Will contain component **prototypes**, **beam tests** and **facilities** with beam

- a muon production and cooling demonstrator is important components
- targets
- hybrid RCS
- ...



Could also feed neutrino facility

Different cooling demonstrator sites are being considered

- CERN, FNAL, ESS are being discussed
- J-PARC also interesting as option

Target could be tested at neutron or muon beam facilities

RCS could be interesting as injectors

Components could be interesting everywhere

Conclusion



- Muon collider is unique opportunity for high-energy, high-luminosity lepton collider
 - Less mature than other options
 - But promises compactness, cost and power efficiency
 - Mainly technological challenges
- Currently two different options considered
 - Goal of 10+ TeV, potential 3 TeV intermediate stage explored
 - Deliverables for next strategies: Project Evaluation Report and R&D Plan
- Collaboration needs US
- Addressing key challenges
 - Very motivated team
 - Synergy with applications for society, e.g. HTS solenoids
 - More funding required for full results by next strategy processes

<http://muoncollider.web.cern.ch>

To join contact

muon.collider.secretariat@cern.ch

Reserve




Staging

Ideally would like full energy right away, but staging could lead to faster implementation

- Substantially less cost for a first stage
- Can make technical compromises
 - e.g. 8 T NbTi magnets would increase collider ring from 4.5 to 6 km and reduce luminosity by 25%
- Can implement improved technology in upgrade


Upgrade adds one more accelerator and new collider ring

- only first collider ring is not being reused



Collider Concept

Fully driven by muon lifetime, otherwise would be easy



Short, intense proton bunch

Ionisation cooling of muon in matter


Acceleration to collision energy

Collision

Protons produce pions which decay into muons
muons are captured

Thanks to MAPS

Key Challenges



Proton complex

- Compressing protons to few bunches

Target

- Target
- Solenoid

Cooling channel

- Channel design
- Solenoids
- RF in magnetic field
- Absorbers
- Integration

RCS


- Beam dynamics
- Ramping magnets
- Power converter
- RF system

Collider ring

- Optics
- Magnets
- Neutrino flux
- Detector background

Protons and Target

MAP target design, K. McDonald, et al.




Large bore, high field solenoid

Nb₃Sn or HTS

Similar to ITER

A. Lechner et al.
L. Bottura et al.




ITER Central Solenoid Model Coil
13 T in 1.7 m (LTS)

2 MW proton beam is OK

Bunching challenge will be
addressed by ESS experts

N. Milas et al. (ESS, Uppsala)

D. Schulte




Shock and power in target:
Simulations of graphite target
indicate 2 MW could be
acceptable

STFC will also study alternatives



Muon Collider, EPP2024, December 2022

Muon Cooling



MAP designs almost achieve 10 TeV goal

- miss factor two for final cooling



MICE Collaboration

Nature vol. 578, p. 53-59 (2020)

Principle demonstrated with no RF
Use of data for benchmarking is still ongoing

Integration/optimisation of design
Integrating improved technologies
Collective effects

C. Rogers et al.
(RAL, CERN)

T. Pieloni et al. (EPFL, CERN)

Cooling Cell Technology



C. Marchand, Alexej Grudiev et al. (CEA, Milano, CERN, Tartu)


RF cavities in magnetic field

MAP demonstrated higher than goal gradient
Improve design based on theoretical understanding
Preparation of **new test stand**, but needs funding

- Test stand at CEA (700 MHz, need funding)
- Test at other frequencies in the UK considered
- Use of CLIC breakdown experiment considered


MuCool demonstrated
>50 MV/m in 5 T

- H_2 -filled copper
- Be end caps



Assessment of realistic goal for highest field solenoids

- MAP demonstrated 30 T
- now magnets aim for 40+ T
- even more can be possible




L. Bottura et al. INFN (Task Leader), CEA, CERN, LNCMI, PSI, SOTON, UNIGE and TWENTE, in collaboration with KEK and US-MDP

Will develop example cooling cell integration

- tight constraints
- additional technologies (**absorbers**, instrumentation,...)
- early preparation of **demonstrator facility**


L. Rossi et al. (INFN, Milano, STFC, CERN), J. Ferreira Somoza et al.



D. Schulte

Muon Collider, EPP2024, December 2022

Acceleration Complex



Core of baseline is sequence of pulsed synchrotron (0.4-11 ms)
Important cost and power consumption

Started

- **Integrated design of RCS**
 - lattice with realistic hardware specifications
 - collective effects
- **Concept of key components**
 - Fast-ramping normal magnets
 - HTS alternative
 - Efficient power converters
 - RF with transient beam loading

A. Chance et al. (CEA)

E. Metral et al. (CERN)

L. Bottura et al. (LNCMI, Darmstadt, Bologna, Twente)

F. Boattini et al.

H. Damerell, F. Batsch, U. van Rienen, A. Grudiev et al. (Rostock, Milano, CERN)

- Alternative FFA S. Machida et al. (RAL)



FNAL 300 T/s HTS magnet



Collider Ring




MAP developed 4.5 km ring for 3 TeV with Nb₃Sn

- magnet specifications in the HL-LHC range

Work progressing on 10 km ring for **10 TeV collider ring**

- around 16 T Nb₃Sn or HTS dipoles
- final focus based on HTS



C. Carli, K. Skoufaris (CERN)

Field choice will be reviewed for cost

Example alternatives:

- a 6 km 3 TeV ring with **NbTi** at 8 T in arcs
- a 15 km 10 TeV ring with HL-LHC performances
- slight reduction in luminosity

15 cm aperture for shielding to ensure magnet lifetime


Need stress managed magnet designs

INFN, Milano, Kyoto, CERN, profit from US

Shielding

A. Lechner
D. Calzolari
(CERN)


Coil




D. Schulte

Muon Collider, EPP2024, December 2022

Alternatives: The LEMMA Scheme



LEMMA scheme (INFN) P. Raimondi et al.





Note: New proposal by C. Curatolo and L. Serafini needs to be looked at

- Uses Bethe-Heitler production with electrons

45 GeV positrons to produce muon pairs
Accumulate muons from several passages

$$e^+ e^- \rightarrow \mu^+ \mu^-$$




Excellent idea, but nature is cruel

Detailed estimates of fundamental limits show that we require a very large positron bunch charge to reach the same luminosity as the proton-based scheme


⇒ **Need same game changing invention**

MICE: Cooling Demonstration

7th February 2015



Nature vol. 578, p. 53-59 (2020)



D. Schulte

More particles at smaller amplitude after absorber is put in place

Principle of ionisation cooling has been demonstrated
Use of data for benchmarking is still ongoing

WEPOPT053

More complete experiment with higher statistics, more than one stage required

Integration of magnets, RF, absorbers, vacuum is engineering challenge

Muon Collider, EPP2024, December 2022

Neutrino Flux

Dense neutrino flux cone can impact environment
Challenge scales with $E \times L$

Goal is to reduce to negligible level, similar to LHC



- 3 TeV, 200 m deep tunnel is about OK

Expand idea of Mokhov, Ginneken to move beam in aperture: move collider ring components, e.g. vertical bending with 1% of main field

- 14 TeV, in 200 m deep tunnel comparable to LHC case with ± 1 mradian
- scales with luminosity toward higher E


Need to study mover system, magnet, connections and impact on beam

Working on different approaches for experimental insertion



Other optimisations are possible (magnetic field, emittance etc.)

Organisation



Thanks



Muon Beam Panel: Daniel Schulte (CERN, chair), Mark Palmer (BNL, co-chair), Tabea Arndt (KIT), Antoine Chance (CEA/IRFU) Jean-Pierre Delahaye (retired), Angeles Faus-Golfe (IN2P3/IJCLab), Simone Gilardoni (CERN), Philippe Lebrun (European Scientific Institute), Ken Long (Imperial College London), Elias Metral (CERN), Nadia Pastrone (INFN-Torino), Lionel Quettier (CEA/IRFU), Magnet Panel link, Tor Raubenheimer (SLAC), Chris Rogers (STFC-RAL), Mike Seidel (EPFL and PSI), Diktys Stratakis (FNAL), Akira Yamamoto (KEK and CERN) **Contributors:** Alexej Grudiev (CERN), Roberto Losito (CERN), Donatella Lucchesi (INFN)

Community conveners: *Radio-Frequency (RF):* Alexej Grudiev (CERN), Jean-Pierre Delahaye (CERN retiree), Derun Li (LBNL), Akira Yamamoto (KEK). *Magnets:* Lionel Quettier (CEA), Toru Ogitsu (KEK), Soren Prestemon (LBNL), Sasha Zlobin (FNAL), Emanuela Barzi (FNAL). *High-Energy Complex (HEC):* Antoine Chance (CEA), J. Scott Berg (BNL), Alex Bogacz (JLAB), Christian Carli (CERN), Angeles Faus-Golfe (IJCLab), Eliana Gianfelice-Wendt (FNAL), Shinji Machida (RAL). *Muon Production and Cooling (MPC):* Chris Rogers (RAL), Marco Calviani (CERN), Chris Densham (RAL), Diktys Stratakis (FNAL), Akira Sato (Osaka University), Katsuya Yonehara (FNAL). *Proton Complex (PC):* Simone Gilardoni (CERN), Hannes Bartosik (CERN), Frank Gerigk (CERN), Natalia Milas (ESS). *Beam Dynamics (BD):* Elias Metral (CERN), Tor Raubenheimer (SLAC and Stanford University), Rob Ryne (LBNL). *Radiation Protection (RP):* Claudia Ahdida (CERN). *Parameters, Power and Cost (PPC):* Daniel Schulte (CERN), Mark Palmer (BNL), Jean-Pierre Delahaye (CERN retiree), Philippe Lebrun (CERN retiree and ESI), Mike Seidel (PSI), Vladimir Shiltsev (FNAL), Jingyu Tang (IHEP), Akira Yamamoto (KEK). *Machine Detector Interface (MDI):* Donatella Lucchesi (University of Padova), Christian Carli (CERN), Anton Lechner (CERN), Nicolai Mokhov (FNAL), Nadia Pastrone (INFN), Sergo R Jindariani (FNAL). *Synergy:* Kenneth Long (Imperial College), Roger Ruber (Uppsala University), Koichiro Shimomura (KEK). *Test Facility (TF):* Roberto Losito (CERN), Alan Bross (FNAL), Tord Ekelof (ESS, Uppsala University).

And the participants to the community meetings and the study

Initial Target Parameters



Target integrated luminosities

$\sqrt{s}$	$\int \mathcal{L} dt$
3 TeV	1 ab ⁻¹
10 TeV	10 ab ⁻¹
14 TeV	20 ab ⁻¹

Note: currently focus on 10 TeV, also explore 3 TeV

- Tentative parameters based on MAP study, might add margins
- Achieve goal in 5 years
- FCC-hh to operate for 25 years
- Aim to have two detectors

Parameter	Unit	3 TeV	10 TeV	14 TeV	CLIC at 3 TeV
L	10 ³⁴ cm ⁻² s ⁻¹	1.8	20	40	2 (6)
N	10 ¹²	2.2	1.8	1.8	
f _r	Hz	5	5	5	
P _{beam}	MW	5.3	14.4	20	28
C	km	4.5	10	14	
	T	7	10.5	10.5	
ε _L	MeV m	7.5	7.5	7.5	
σ _E / E	%	0.1	0.1	0.1	
σ _z	mm	5	1.5	1.07	
β	mm	5	1.5	1.07	
ε	μm	25	25	25	
σ _{x,y}	μm	3.0	0.9	0.63	

Roadmap



In **aspirational scenario** can make **informed decisions**:

Three main deliverables are foreseen:

- a **Project Evaluation Report** for the next ESPPU will contain an assessment of whether the 10 TeV muon collider is a promising option and identify the required compromises to realise a 3 TeV option by 2045. In particular the questions below would be addressed.
 - What is a realistic luminosity target?
 - What are the background conditions in the detector?
 - Can one consider implementing such a collider at CERN or other sites, and can it have one or two detectors?
 - What are the key performance specifications of the components and what is the maturity of the technologies?
 - What are the cost drivers and what is the cost scale of such a collider?
 - What are the power drivers and what is the power consumption scale of the collider?
 - What are the key risks of the project?
- an **R&D Plan** that describes an R&D path towards the collider;
- an **Interim Report** by the end of 2023 that documents progress and allows the wider community to update their view of the concept and to give feedback to the collaboration.

R&D Plan



The R&D plan will describe the R&D path toward the collider, in particular during the CDR phase, and will comprise the elements below.

- An integrated concept of a muon cooling cell that will allow construction and testing of this key novel component.
- A concept of the facility to provide the muon beam to test the cells.
- An evaluation of whether this facility can be installed at CERN or another site.
- A description of other R&D efforts required during the CDR phase including other demonstrators.

This R&D plan will allow the community to understand the technically limited timeline for the muoncollider development after the next ESPPU.

Minimal Scenario



Will allow **partially informed decisions**

- No conceptual design of neutrino flux and alignment system
- No alternative superconducting fast-ramping magnet system
- Several collider systems would (almost) not be covered, in particular
 - the linacs
 - the target complex
 - the proton complex
 - engineering considerations of the muon cooling cells
 - alternative designs for the final cooling system, acceleration, collider ring
- No RF test stand would be constructed for the muon cooling accelerating cavities
- No conceptual design of a muon cooling cell for the test programme
- No conceptual design of a muon cooling demonstrator facility
- No concept of RF power sources
- No tests/models to develop solenoid technology.

Schedule




Fig. 5.4: Overall timeline for the R&D programme.

2023

2025

2025-2027

Interim Report to gauge progress
Initial baseline defined

Assessment Report

R&D plan will be refined


Test Facility Dimensions

Target

+ horn (1st phase) /
+ superconducting solenoid (2nd phase)

**Collimation and upstream
diagnostics area, 10 m**

**Downstream
diagnostics area, 5 m**




Momentum selection chicane, 10 m

Cooling area, 50 m

Look for an existing proton beam with
significant power

Different sites are being considered



- CERN, FNAL, ESS are being discussed
- J-PARC also interesting as option



Possible CERN Locations

Consider nTOF-like beam from PS for cooling experiment:

- 1 pulse of 10^{13} p at 20 GeV per 1.2 s, i.e. 27 kW
 - maybe O(100kW) possible
- If SPL were, installed could use its beam, e.g. 5 GeV, 4 MW



MoC and Design Study Partners



IEIO	CERN
FR	CEA-IRFU
	CNRS-LNCMI
DE	DESY
	Technical University of Darmstadt
	University of Rostock
	KIT
IT	INFN
	INFN, Univ., Polit. Torino
	INFN, Univ. Milano
	INFN, Univ. Padova
	INFN, Univ. Pavia
	INFN, Univ. Bologna
	INFN Trieste
	INFN, Univ. Bari
	INFN, Univ. Roma 1
	ENEA

UK	RAL
	UK Research and Innovation
	University of Lancaster
	University of Southampton
	University of Strathclyde
	University of Sussex
	Imperial College
	Royal Holloway
	University of Huddersfield
	University of Oxford
	University of Warwick
	University of Durham
SE	ESS
	University of Uppsala
PT	LIP
NL	University of Twente

FI	Tampere University
US	Iowa State University
	Wisconsin-Madison
	Pittsburg University
	BNL
China	Sun Yat-sen University
	IHEP
	Peking University
EST	Tartu University
LAT	Riga Technical Univers.
AU	HEPHY
	TU Wien
ES	IBM
CH	PSI
	University of Geneva
	EPFL
BE	Louvain

IT	INFN Frascati
	INFN, Univ. Ferrara
	INFN, Univ. Roma 3
	INFN Legnaro
	INFN, Univ. Milano Bicocca
	INFN Genova
	INFN Laboratori del Sud
	INFN Napoli
US	FNAL
	LBL
	JLAB
	Chicago
Japan	Akira Yamamoto
	Akira Sato
	Toru Ogitsu

D. Schulte

Muon Collider, EPP2024, December 2022

Coordination Committee Members



Physics	Andrea Wulzer
Detector and MDI	Donatella Lucchesi

Protons	Natalia Milas
Muon production and cooling	Chris Rogers
Muon acceleration	Antoine Chance
Collider	Christian Carli

Magnets	Luca Bottura
RF	Alexej Grudiev, Claude Marchand
Beam-matter int. target systems	Anton Lechner
Collective effects	Elias Metral

Cooling cell design	Lucio Rossi
Demonstrator	Roberto Losito

US (detector)	Sergo Jindariani
US (accelerator)	Mark Palmer
Asia (China)	Jingyu Tang
Asia (Japan)	tbd

A strengthening on the physics and detector side is planned

Key Technologies



Magnets

- Superconducting solenoids for target and cooling profit from developments for society
 - target solenoid comparable to ITER central solenoid fusion
 - 6D cooling solenoids similar and wind power generators, motors
 - final cooling solenoids synergetic with high-field research, NMR
- Collider ring magnets
 - profit from developments for other colliders FCC-hh, stress-managed magnets
- Fast-ramping normal-conducting magnet system
 - HTS alternative, power converter

Key Technologies, cont.



RF systems

- Normal-conducting cooling cavities in magnetic field
 - profit from CLIC work
- Superconducting accelerator RF
 - profit from ILC, ...
- Efficient power sources
 - profit from CLIC work

Beam-matter interaction

- Proton target
- Cooling absorbers
- Shielding (accelerator and detector)

Mechanical system

- Neutrino flux mitigation system
- Muon cooling cell integration