
LIGO Discovery of a Binary Black-Hole Merger



Vicky Kalogera


Dept of Physics & Astronomy
Center for Interdisciplinary Exploration and Research
in Astrophysics (CIERA)

for the LIGO-Virgo Collaborations












LIGO Timeline





ZLIGO

LIGO Scientific Collaboration









































































Northwestern

















UNIVERSITYOF



















UNIVERSITY of WISCONSIN MILWAUKEE

























Leibniz Universität Hannover










LIGO Team

- Astrophysics of LIGO sources
 - computational modeling of compact object binaries
 - predictions for LIGO observations
 - interpretation of observed systems

- LIGO data analysis
 - advanced method development
 - data characterization
 - source parameter estimation

Binary Inspiral Source of Gravitational Waves










Binary Inspiral Source of Gravitational Waves











Binary Inspiral Source of Gravitational Waves











Two LIGO detectors








Initial LIGO

2001 - 2006


2006 - 2010

Laser Interferometer






Laser Interferometer





Initial LIGO





Advanced LIGO: Plan for Observing Runs

You are here


NS-NS Reach in Mpc

Second generation interferometers to begin science operations:

- Advanced LIGO (2 interferometers) 2015
- Advanced Virgo (1 interferometer) 2016

Approximate run schedule:

- Advanced LIGO:
 - ~ 3 month run in 2015,
 - ~ 6 month run in 2016-17
 - ~ 9 month run in 2017-18
- · Advanced Virgo:
 - ~ 6 month run in 2016-17
 - ~ 9 month run in 2017-18
- Modification of run schedules is likely as we learn more about the instruments






Advanced LIGO

Key Detector Upgrades

- Increased Laser Power
- Bigger Mirror Masses
- Better Mirror Coatings
- Improved Seismic Isolation











 $S/N \sim 20$

peak strain: 1e-21









signal and source

$$\mathcal{M} = \frac{(m_1 m_2)^{3/5}}{(m_1 + m_2)^{1/5}} = \frac{c^3}{G} \left[\frac{5}{96} \pi^{-8/3} f^{-11/3} \dot{f} \right]^{3/5}.$$

			ı	
	observed by	LIGO L1, H1	duration from 30 Hz	~ 200 ms
	source type	black hole (BH) binary	# cycles from 30 Hz	~10
	date	14 Sept 2015	peak GW strain	1 x 10 ⁻²¹
	time	09:50:45 UTC	peak displacement of	10 000 fm
	likely distance	0.75 to 1.9 Gly	interferometers arms	±0.002 fm
		230 to 570 Mpc	frequency/wavelength	150 Hz, 2000 km
	redshift	0.054 to 0.136	at peak GW strain	
S	ignal-to-noise ratio	24	peak speed of BHs	~ 0.6 c
	false alarm prob.	less than 1 in 5 million	peak GW luminosity	3.6 x 10 ⁵⁶ erg s ⁻¹
	false alarm rate		radiated GW energy	2.5-3.5 M⊙
	Source Mas	1 in 200,000 yr	remnant ringdown fre	q. ~ 250 Hz
	THE STATE OF THE S		remnant damping tim	ne ~ 4 ms
	total mass	65		
	chirpmass	28	remnant size, area	180 km, 3.5 x 10 ⁵ km ²
	primary BH	32 to 41	consistent with	passes all tests
	secondary BH	25 to 33	general relativity?	performed
	remnant BH	62	graviton mass bound	< 1.2 x 10 ⁻²² eV
	mass ratio	0.6 to 1	coalescence rate	2 to 400 Gpc ⁻³ yr ⁻¹
	primary BH spin	< 0.7	online trigger latency	~ 3 min
5	secondary BH spin	< 0.9	# offline analysis pipeli	nes 5
	remnant BH spin	0.7	CDU I	~ 50 million (=20,000
:	signal arrival time	arrived in L1 7 ms	CPU hours consumed	PCs run for 100 days)
	delay	before H1	papers on Feb 11, 2016	13
	likely sky position	Southern Hemisphere	1, 2010	A A A A A A A A A A A A A A A A A A A
M	likely orientation resolved to	face-on/off ~600 sq. deg.	# researchers	~1000, 80 institutions in 15 countries
		~ W \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \		V V V V V V V V V V V V V V V V V V V



signal and source

$$\mathcal{M} = \frac{(m_1 m_2)^{3/5}}{(m_1 + m_2)^{1/5}} = \frac{c^3}{G} \left[\frac{5}{96} \pi^{-8/3} f^{-11/3} \dot{f} \right]^{3/5}.$$

			1	
	observed by	LIGO L1, H1	duration from 30 Hz	~ 200 ms
	source type	black hole (BH) binary	# cycles from 30 Hz	~10
	date	14 Sept 2015	peak GW strain	1 x 10 ⁻²¹
	time	09:50:45 UTC	peak displacement of	. 0 000 (
	likely distance	0.75 to 1.9 Gly 230 to 570 Mpc	interferometers arms frequency/wavelength	±0.002 fm 150 Hz, 2000 km
	redshift	0.054 to 0.136	at peak GW strain	130 112, 2000 KIII
	signal-to-noise ratio	24	peak speed of BHs	~ 0.6 c 3.6 x 10 ⁵⁶ erg s ⁻¹
	false alarm prob.	less than 1 in 5 million	peak GW luminosity radiated GW energy	2.5-3.5 M⊙
	false alarm rate	1 in 200,000 yr		250 II-
	Source Mass	ses M⊙	remnant ringdown fre	
	total mass	65	remnant damping tim	ne ~ 4 ms
	chirpmass	28	remnant size, area	180 km, 3.5 x 10 ⁵ km ²
5	primary BH	32 to 41	consistent with	passes all tests
	secondary BH	25 to 33	general relativity?	performed
	remnant BH	62	graviton mass bound	< 1.2 x 10 ⁻²² eV
	mass ratio	0.6 to 1	coalescence rate	2 to 400 Gpc ⁻³ yr ⁻¹
	primary BH spin	< 0.7	online trigger latency	~ 3 min
	secondary BH spin	< 0.9	# offline analysis pipeli	nes 5
	remnant BH spin	0.7	CDU have as reversed	~ 50 million (=20,000
	signal arrival time	arrived in L1 7 ms	CPU hours consumed	PCs run for 100 days)
	delay	before H1	papers on Feb 11, 2016	13
	likely sky position	Southern Hemisphere	The stand of the standard	1000 80 institutions
	likely orientation resolved to	face-on/off ~600 sq. deg.	# researchers	~1000, 80 institutions in 15 countries
		 	+	<u> </u>



signal and source

$$\mathcal{M} = \frac{(m_1 m_2)^{3/5}}{(m_1 + m_2)^{1/5}} = \frac{c^3}{G} \left[\frac{5}{96} \pi^{-8/3} f^{-11/3} \dot{f} \right]^{3/5}.$$

observed by	LIGO L1, H1	duration from 30 Hz ~ 200 ms
source type	black hole (BH) binary	# cycles from 30 Hz ~10
date	14 Sept 2015	peak GW strain 1 x 10 ⁻²¹
time	09:50:45 UTC	nook dienlocoment of
likely distance	0.75 to 1.9 Gly 230 to 570 Mpc	peak displacement of ±0.002 fm interferometers arms frequency/wavelength 150 Hz, 2000 km
redshift	0.054 to 0.136	at peak GW strain
signal-to-noise rat	io 24	peak speed of BHs ~ 0.6 c
false alarm prob.	less than 1 in 5 million	peak GW luminosity 3.6 x 10 ⁵⁶ erg s ⁻¹ radiated GW energy 2.5-3.5 M☉
false alarm rate	1 in 200,000 yr	
Source M	lasses M⊙	remnant ringdown freq. ~ 250 Hz
total mass	65	remnant damping time ~ 4 ms
chirpmass	28	remnant size, area 180 km, 3.5 x 10 ⁵ km ²
primary BH	32 to 41	consistent with passes all tests
secondary BH	25 to 33	general relativity? pe <mark>rfo</mark> rmed
remnant BH	62	graviton mass bound < 1.2 x 10 ⁻²² eV
mass ratio	0.6 to 1	coalescence rate 2 to 400 Gpc ⁻³ yr ⁻¹
primary BH spin		online trigger latency ~ 3 min
secondary BH spi	n < 0.9	# offline analysis pipelines 5
remnant BH spin signal arrival time	e arrived in L1 7 ms	CPU hours consumed CPU hours consumed PCs run for 100 days)
delay likely sky positior	before H1 Southern Hemisphere	papers on Feb 11, 2016 13
likely orientation resolved to		# researchers in 15 countries



signal and source

$$\mathcal{M} = \frac{(m_1 m_2)^{3/5}}{(m_1 + m_2)^{1/5}} = \frac{c^3}{G} \left[\frac{5}{96} \pi^{-8/3} f^{-11/3} \dot{f} \right]^{3/5}.$$

	observed by	LIGO L1, H1	duration from 30 Hz	~ 200 ms
	source type	black hole (BH) binary	# cycles from 30 Hz	~10
	date	14 Sept 2015	peak GW strain	1 x 10 ⁻²¹
	time	09:50:45 UTC	nook displacement of	
	likely distance	0.75 to 1.9 Gly 230 to 570 Mpc	peak displacement of interferometers arms frequency/wavelength	±0.002 fm
	redshift	0.054 to 0.136	at peak GW strain	150 Hz, 2000 km
9	signal-to-noise ratio	24	peak speed of BHs	~ 0.6 c
	false alarm prob.	less than 1 in 5 million	peak GW luminosity radiated GW energy	3.6 x 10 ⁵⁶ erg s ⁻¹ 2.5-3.5 M⊙
	false alarm rate	1 in 200,000 yr		
	Source Mas	sses Mo	remnant ringdown fred	
	total mass	65	remnant damping tim	e ~ 4 ms
	chirpmass	28	remnant size, area	180 km, 3.5 x 10 ⁵ km ²
5	primary BH	32 to 41	consistent with	passes all tests
	secondary BH	25 to 33	general relativity?	performed
	remnant BH	62	graviton mass bound	< 1.2 x 10 ⁻²² eV
	mass ratio	0.6 to 1	coalescence rate	2 to 400 Gpc ⁻³ yr ⁻¹
	primary BH spin	< 0.7	online trigger latency	~ 3 min
	secondary BH spin	< 0.9	# offline analysis pipelir	nes 5
1	remnant BH spin	0.7		~ 50 million (=20,000
	signal arrival time	arrived in L1 7 ms	CPU hours consumed	PCs run for 100 days)
	delay	before H1	papers on Feb 11, 2016	13
A	likely sky position	Southern Hemisphere		MA MANA
1	likely orientation resolved to	face-on/off ~600 sq. deg.	# researchers	~1000, 80 institutions in 15 countries



signal and source

$$\mathcal{M} = \frac{(m_1 m_2)^{3/5}}{(m_1 + m_2)^{1/5}} = \frac{c^3}{G} \left[\frac{5}{96} \pi^{-8/3} f^{-11/3} \dot{f} \right]^{3/5}.$$

		ı	
observed by	LIGO L1, H1	duration from 30 Hz	~ 200 ms
source type	black hole (BH) binary	# cycles from 30 Hz	~10
date	14 Sept 2015	peak GW strain	1 x 10 ⁻²¹
time	09:50:45 UTC	peak displacement of	
likely distance	0.75 to 1.9 Gly	interferometers arms	±0.002 fm
likely distance	230 to 570 Mpc	frequency/wavelength	150 Hz, 2000 km
redshift	0.054 to 0.136	at peak GW strain	130 112, 2000 KIII
signal-to-noise ratio	24	peak speed of BHs	~ 0.6 c
	less than 1 in 5 million	peak GW luminosity	3.6 x 10 ⁵⁶ erg s ⁻¹
false alarm prob.	less than 1 in 5 million	radiated GW energy	2.5-3.5 M⊙
false alarm rate	1 in 200,000 yr	romport ring down fro	g. ~ 250 Hz
Source Mas	sses M⊙	remnant ringdown fre	-
total mass	65	remnant damping tin	
chirpmass	28	remnant size, area	180 km, 3.5 x 10 ⁵ km ²
primary BH	32 to 41	consistent with	passes all tests
secondary BH	25 to 33	general relativity?	pe <mark>rfo</mark> rmed
remnant BH	62	graviton mass bound	< 1.2 x 10 ⁻²² eV
mass ratio	0.6 to 1	coalescence rate	2 to 400 Gpc ⁻³ yr ⁻¹
primary BH spin	< 0.7	online trigger latency	~ 3 min
secondary BH spin	< 0.9	# offline analysis pipeli	N /
remnant BH spin	0.7		~ 50 million (=20,000
signal arrival time	arrived in L1 7 ms	CPU hours consumed	PCs run for 100 days)
delay	before H1	papers on Feb 11, 2016	13
likely sky position	Southern Hemisphere	papers on Feb 11, 2010	A AA A
likely orientation resolved to	face-on/off ~600 sq. deg.	# researchers	~1000, 80 institutions in 15 countries



signal and source

$$\mathcal{M} = \frac{(m_1 m_2)^{3/5}}{(m_1 + m_2)^{1/5}} = \frac{c^3}{G} \left[\frac{5}{96} \pi^{-8/3} f^{-11/3} \dot{f} \right]^{3/5}.$$

			I	
	observed by	LIGO L1, H1	duration from 30 Hz	~ 200 ms
	source type	black hole (BH) binary	# cycles from 30 Hz	~10
	date	14 Sept 2015	peak GW strain	1 x 10 ⁻²¹
I	time	09:50:45 UTC	peak displacement of	
	likely distance	0.75 to 1.9 Gly	interferometers arms	±0.002 fm
	intery distance	230 to 570 Mpc	frequency/wavelength	150 Hz, 2000 km
	redshift	0.054 to 0.136	at peak GW strain	
7	signal-to-noise ratio	24	peak speed of BHs	~ 0.6 c
	false alarm prob.	less than 1 in 5 million	peak GW luminosity	3.6 x 10 ⁵⁶ erg s ⁻¹
			radiated GW energy	2.5-3.5 M⊙
_	false alarm rate	1 in 200,000 yr	remnant ringdown free	g. ~ 250 Hz
_	Source Mas	sses M⊙		
	total mass	65	remnant damping tim	
	chirpmass	28	remnant size, area	180 km, 3.5 x 10 ⁵ km ²
	primary BH	32 to 41	consistent with	passes all tests
	secondary BH	25 to 33	general relativity?	pe <mark>rfo</mark> rmed
	remnant BH	62	graviton mass bound	< 1.2 x 10 ⁻²² eV
	mass ratio	0.6 to 1	coalescence rate	2 to 400 Gpc ⁻³ yr ⁻¹
	primary BH spin	< 0.7	online trigger latency	~ 3 min
	secondary BH spin	< 0.9	# offline analysis pipelii	nes 5
	remnant BH spin	0.7		~ 50 million (=20,000
	signal arrival time	arrived in L1 7 ms	CPU hours consumed	PCs run for 100 days)
	delay	before H1	papers on Feb 11, 2016	13
	likely sky position	Southern Hemisphere	Papers Off 1 60 11, 2010	A AM MAIN
	likely orientation resolved to	face-on/off ~600 sq. deg.	# researchers	~1000, 80 institutions in 15 countries
			++	



signal and source

$$\mathcal{M} = \frac{(m_1 m_2)^{3/5}}{(m_1 + m_2)^{1/5}} = \frac{c^3}{G} \left[\frac{5}{96} \pi^{-8/3} f^{-11/3} \dot{f} \right]^{3/5}.$$

	observed by	LIGO L1, H1	duration from 30 Hz ~ 200 ms
	source type	black hole (BH) binary	# cycles from 30 Hz ~10
	date	14 Sept 2015	peak GW strain 1 x 10 ⁻²¹
	time	09:50:45 UTC	peak displacement of ±0.002 fm
	likely distance	0.75 to 1.9 Gly	interferometers arms
		230 to 570 Mpc	frequency/wavelength 150 Hz, 2000 km
	redshift	0.054 to 0.136	at peak GW strain
	signal-to-noise ratio	24	peak speed of BHs ~ 0.6 c
	false alarm prob.	less than 1 in 5 million	peak GW luminosity 3.6 x 10 ⁵⁶ erg s ⁻¹
	false alarm rate	1 in 200,000 yr	radiated GW energy 2.5-3.5 M⊙
_	Source Ma		remnant ringdown freq. ~ 250 Hz
_			remnant damping time ~ 4 ms
	total mass	65	
	chirpmass	28	remnant size, area 180 km, 3.5 x 10 ⁵ km ²
5	primary BH	32 to 41	consistent with passes all tests
	secondary BH	25 to 33	general relativity? performed
	remnant BH	62	graviton mass bound < 1.2 x 10 ⁻²² eV
	mass ratio	0.6 to 1	coalescence rate 2 to 400 Gpc ⁻³ yr ⁻¹
	primary BH spin	< 0.7	online trigger latency ~ 3 min
	secondary BH spin	< 0.9	
\	remnant BH spin	0.7	, , , , , , , , , , , , , , , , , , ,
Г	signal arrival time	arrived in L1 7 ms	CPU hours consumed ~ 50 million (=20,000 PCs run for 100 days)
	delay	before H1	
	likely sky position	Southern Hemisphere	papers on Feb 11, 2016 13
	likely orientation resolved to	face-on/off ~600 sq. deg.	# researchers ~1000, 80 institutions in 15 countries



signal and source

$$\mathcal{M} = \frac{(m_1 m_2)^{3/5}}{(m_1 + m_2)^{1/5}} = \frac{c^3}{G} \left[\frac{5}{96} \pi^{-8/3} f^{-11/3} \dot{f} \right]^{3/5}.$$

		1	
observed by	LIGO L1, H1	duration from 30 Hz	~ 200 ms
source type	black hole (BH) binary	# cycles from 30 Hz	~10
date	14 Sept 2015	peak GW strain	1 x 10 ⁻²¹
time	09:50:45 UTC	peak displacement of	.0.000 (
likely distance	0.75 to 1.9 Gly	interferometers arms	±0.002 fm
	230 to 570 Mpc	frequency/wavelength	150 Hz, 2000 km
redshift	0.054 to 0.136	at peak GW strain	
signal-to-noise ratio	24	peak speed of BHs	~ 0.6 c
false alarm prob.	less than 1 in 5 million	peak GW luminosity	3.6 x 10 ⁵⁶ erg s ⁻¹
false alarm rate	1 in 200,000 yr	radiated GW energy	2.5-3.5 M⊙
Source Mas		remnant ringdown fre	q. ~ 250 Hz
		remnant damping tim	ne ~ 4 ms
total mass	65		
chirpmass	28	remnant size, area	180 km, 3.5 x 10 ⁵ km ²
primary BH	32 to 41	consistent with	passes all tests
secondary BH	25 to 33	general relativity?	pe <mark>rfo</mark> rmed
remnant BH	62	graviton mass bound	< 1.2 x 10 ⁻²² eV
mass ratio	0.6 to 1	coalescence rate	2 to 400 Gpc ⁻³ yr ⁻¹
primary BH spin	< 0.7	online trigger latency	~ 3 min
secondary BH spin	< 0.9	# offline analysis pipeli	nes 5
remnant BH spin	0.7		~ 50 million (=20,000
signal arrival time	arrived in L1 7 ms	CPU hours consumed	PCs run for 100 days)
delay	before H1	papers on Feb 11, 2016	13
likely sky position	Southern Hemisphere	Papers 0111 CD 11, 2010	
likely orientation resolved to	face-on/off ~600 sq. deg.	# researchers	~1000, 80 institutions in 15 countries



signal and source

$$\mathcal{M} = \frac{(m_1 m_2)^{3/5}}{(m_1 + m_2)^{1/5}} = \frac{c^3}{G} \left[\frac{5}{96} \pi^{-8/3} f^{-11/3} \dot{f} \right]^{3/5}.$$

		I	
observed by	LIGO L1, H1	duration from 30 Hz	~ 200 ms
source type	black hole (BH) binary	# cycles from 30 Hz	~10
date	14 Sept 2015	peak GW strain	1 x 10 ⁻²¹
time	09:50:45 UTC	peak displacement of	.0.002.0
likely distance	0.75 to 1.9 Gly	interferometers arms	±0.002 fm
	230 to 570 Mpc	frequency/wavelength	150 Hz, 2000 km
redshift	0.054 to 0.136	at peak GW strain	
signal-to-noise ratio	24	peak speed of BHs	~ 0.6 c
false alarm prob.	less than 1 in 5 million	peak GW luminosity	3.6 x 10 ⁵⁶ erg s ⁻¹
false alarm rate		radiated GW energy	2.5-3.5 M⊙
Source Mas	1 in 200,000 yr sses M⊙	remnant ringdown fre	q. ~ 250 Hz
		remnant damping tim	ne ~ 4 ms
total mass	65	remnant size, area	180 km, 3.5 x 10 ⁵ km ²
chirpmass	28	consistent with	passes all tests
primary BH secondary BH	32 to 41 25 to 33	general relativity?	performed
remnant BH	62	graviton mass bound	< 1.2 x 10 ⁻²² eV
mass ratio	0.6 to 1	coalescence rate	2 to 400 Gpc ⁻³ yr ⁻¹
primary BH spin	< 0.7	online trigger latency	~ 3 min
secondary BH spin	< 0.9	# offline analysis pipeli	
remnant BH spin	0.7		~ 50 million (=20,000
signal arrival time	arrived in L1 7 ms	CPU hours consumed	PCs run for 100 days)
delay	before H1	papers on Feb 11, 2016	W I I
likely sky position	Southern Hemisphere	papers off reb 11, 2010	
likely orientation	face-on/off	# researchers	~1000, 80 institutions
resolved to	~600 sq. deg.	In lead with a pale, Mys.	in 15 countries
	V		V W VI V VI




signal and source

$$\mathcal{M} = \frac{(m_1 m_2)^{3/5}}{(m_1 + m_2)^{1/5}} = \frac{c^3}{G} \left[\frac{5}{96} \pi^{-8/3} f^{-11/3} \dot{f} \right]^{3/5}.$$

			ı	
	observed by	LIGO L1, H1	duration from 30 Hz	~ 200 ms
	source type	black hole (BH) binary	# cycles from 30 Hz	~10
	date	14 Sept 2015	peak GW strain	1 x 10 ⁻²¹
	time	09:50:45 UTC	peak displacement of	10 000 fm
	likely distance	0.75 to 1.9 Gly	interferometers arms	±0.002 fm
		230 to 570 Mpc	frequency/wavelength	150 Hz, 2000 km
	redshift	0.054 to 0.136	at peak GW strain	
S	ignal-to-noise ratio	24	peak speed of BHs	~ 0.6 c
	false alarm prob.	less than 1 in 5 million	peak GW luminosity	3.6 x 10 ⁵⁶ erg s ⁻¹
	false alarm rate		radiated GW energy	2.5-3.5 M⊙
	Source Mas	1 in 200,000 yr	remnant ringdown fre	q. ~ 250 Hz
	THE STATE OF THE S		remnant damping tim	ne ~ 4 ms
	total mass	65		
	chirpmass	28	remnant size, area	180 km, 3.5 x 10 ⁵ km ²
	primary BH	32 to 41	consistent with	passes all tests
	secondary BH	25 to 33	general relativity?	performed
	remnant BH	62	graviton mass bound	< 1.2 x 10 ⁻²² eV
	mass ratio	0.6 to 1	coalescence rate	2 to 400 Gpc ⁻³ yr ⁻¹
	primary BH spin	< 0.7	online trigger latency	~ 3 min
5	secondary BH spin	< 0.9	# offline analysis pipeli	nes 5
	remnant BH spin	0.7		~ 50 million (=20,000
:	signal arrival time	arrived in L1 7 ms	CPU hours consumed	PCs run for 100 days)
	delay	before H1	papers on Feb 11, 2016	13
	likely sky position	Southern Hemisphere	1, 2010	A A A A A A A A A A A A A A A A A A A
M	likely orientation resolved to	face-on/off ~600 sq. deg.	# researchers	~1000, 80 institutions in 15 countries
		- W / W -		V V V V V V V V V V V V V V V V V V V



credit: C. Pankow





GW150914 Papers

Discovery paper

"Observation of Gravitational Waves from a Binary Black Hole Merger" before Feb 11 — Published in PRL 116, 061102 (2016)

Companion papers:

"Astrophysical Implications of the Binary Black-Hole Merger GW150914" before Feb 11 — Published in ApJL 818, L22 (2016)

"Observing gravitational-wave transient GW150914 with minimal assumptions"

"GW150914: First results from the search for binary black hole coalescence with Advanced LIGO"

"Properties of the binary black hole merger GW150914"

"The Rate of Binary Black Hole Mergers Inferred from Advanced LIGO Observations Surrounding GW150914"



GW150914 Papers

Companion papers:

"Tests of general relativity with GW150914"

"GW150914: Implications for the stochastic gravitational-wave background from binary black holes"

"Calibration of the Advanced LIGO detectors for the discovery of the binary black-hole merger GW150914"

"Characterization of transient noise in Advanced LIGO relevant to gravitational wave signal GW150914"

"High-energy Neutrino follow-up search of Gravitational Wave Event GW150914 with IceCube and ANTARES"

"GW150914: The Advanced LIGO Detectors in the Era of First Discoveries"

"Localization and broadband follow-up of the gravitational-wave transient GW150914"



GW150914 Analyses Results



LIGO Searches

39 days of LIGO observations: Sep 12 - Oct 20 16 days of coincident time


Data searched with 5 independent pipelines:

- 3 burst pipelines
- 2 binary-coalescence pipelines all give consistent results!!!

• Key analysis method: "time slides"




LIGO Search for Compact Binary Coalescence





LIGO Search for Compact Binary Coalescence





Physical Parameter Estimation


Bayesian stochastic sampling of 12D parameter space:

- 2 different samplers
- 2 different binary-coalescence waveforms
- 1 generic "waveform"

all give consistent results!!!



GW150914: Masses




individual masses


The LVC, PRL, arxiv/1602.03840



final mass and spin (from NR fits)



GW150914: Spins




spins aligned with orb. ang.mom. are constrained to be small


in-plane spins are unconstrained



GW150914: distance - inclination



~600 square degrees





The LVC, PRL, arxiv/1602.03840



GW150914: sky map

GW150914: Binary BH merger rate






current estimate: 2 - 400 per Gpc³ per yr model predictions: 0 - 1,000 per Gpc³ per yr

rates below ~1 per Gpc³ per yr are excluded




GW150914 EM Follow-up



All upper limits except potential Fermi GBM counterpart?





GW150914 Astrophysics




- First Binary BH system
- Heaviest stellar-mass Black Holes (>~ 25 Msun)

Belczynski et al. 2010

Spera et al. 201:

- First Binary BH system
- Heaviest stellar-mass Black Holes (>~ 25 Msun)



Z < 1/2 solar



BBH Formation

Isolated Binaries

low - Z to PopIII

rapid rotation

Dense Clusters

globular clusters

young clusters

galactic centers



Binary Population Synthesis: some bibliography

Kornilov & Lipunov 1984 Dewey & Cordes 1987

including BBH:

Lipunov et al. 1997

Bethe & Brown 1998

DeDonder & Vanbeveren 1998, 2004

Portegies Zwart & Yungelson 1998

Belczynski & Bulik 1999

Bloom et al. 1999

Fryer et al. 1999

Grishchuk et al. 2001

Nelemans et al. 2001

Belczynski, VK, & Bulik 2002

Voss & Tauris 2003

O'Shaughnessy, 2005

Belczynski et al. 2008


O'Shaughnessy, VK, Belczynski 2010

including heavy BBH:

Belczynski et al. 2010 Fryer et al. 2012 Dominik et al. 2012, 2013, 2015 Belczynski et al. 2015 Spera et al. 2015

BBH Formation from Isolated Binaries

BH masses &
BBH rates:
consistent



stable mass transfer

common envelope

low BH kicks

Belczynski et al. 2016

BBH from Dense Stellar Environments

Globular Clusters some bibliography ...

> Just BH dynamics

Sigurdsson & Hernquist 1993 Portegies Zwart & McMillan 2000 Gultekin et al. 2004, 2006 Kocsis, Gaspar, & Marka 2006 Banerjee et al. 2010 Bae, Kim, & Lee 2014

> Binary Evolution & Semi-Dynamics

O'Leary, ..., O'Shaughnessy 2006 O'Leary, O'Shaughnessy, & Rasio 2007 Sadowski et al. 2008

> Dynamics & Binary Evolution

Downing et al. 2010 Downing et al. 2011

Pattabiraman, ..., VK, Rasio 2013 Morscher et al. 2013 Morscher et al. 2015 Antonini, ..., VK, Rasio 2015

Galactic Centers


> Just BH dynamics Miller & Lauburg 2009 O'Leary et al. 2009 cf. Tsang 2013 Kocsis & Levin 2012

Young Star Clusters


Goswami, Kiel, & Rasio 2014 Ziosi, ..., Branchesi, & Tormen 2014

BBH from globular clusters

Rodriguez, et al. 2016a



BBH Formation in Globular Clusters



BH masses &
BBH rates:
consistent


Optimal host globular cluster:

typical mass: ~5e5 solar masses larg-ish virial radius: 2pc


Predicted rates:

2 - 20 per Gpc³ per yr

Rodriguez et al. 2016b



BBH Formation in Globular Clusters



BH masses &
BBH rates:
consistent


Optimal host globular cluster:

typical mass: ~5e5 solar masses larg-ish virial radius: 2pc

Predicted rates:

2 - 20 per Gpc³ per yr

Rodriguez et al. 2016b



GW150914 and BBH Formation


- heavy black holes exist in merging binaries
- their formation requires an origin from lowmetallicity environments (1/2 solar or less)
- such BBH can form both isolated binaries and dynamical processes
- both formation channels produce merger rates compatible with the event
- the low end of rate predictions (<1 per Gpc³ per yr) is excluded



Ready for the Era of Gravitational-Wave Astrophysics?



Ready for the Era of Gravitational-Wave Astrophysics?



Advanced LIGO: Plan for Observing Runs

You are here


NS-NS Reach in Mpc

Second generation interferometers to begin science operations:

- Advanced LIGO (2 interferometers) 2015
- Advanced Virgo (1 interferometer) 2016

Approximate run schedule:

- Advanced LIGO:
 - ~ 3 month run in 2015,
 - ~ 6 month run in 2016-17
 - ~ 9 month run in 2017-18
- · Advanced Virgo:
 - ~ 6 month run in 2016-17
 - ~ 9 month run in 2017-18
- Modification of run schedules is likely as we learn more about the instruments





Ready for the Era of Gravitational-Wave Astrophysics?


More BBH detections to come ...

Reveal underlying BBH mass distribution

Quantitative model constraints


The LVC, ApJL, arxiv/1602.03842








The advanced GW detector network




The Gravitational Wave Spectrum



Slide: Matt Evans (MIT)

What does GW150914 imply for LISA?

Sesana 2016, PRL, submitted, arXiv:1602.06951



up to ~thousands of BBH resolved by eLISA

of these BBH, up to ~hundreds of mergers will merge within the aLIGO band within 10yr

GW150914: our first Binary BH merger


-0.76s



GW150914: our first Binary BH merger

-0.76s






BBH Formation from Isolated Binaries Common-Envelope & Metallicity Influence

Dominik, Belczynski, ..., Holz, ..., Bulik, Mandel, O'Shaughnessy 2012

CE before & on Giant Branch CE only Giant Branch




solar metallicity

0.1xsolar metallicity

BBH Formation from Isolated Binaries Common-Envelope & Metallicity Influence

Dominik, Belczynski, ..., Holz, ..., Bulik, Mandel, O'Shaughnessy 2012


CE before & on Giant Branch CE only Giant Branch



solar metallicity

0.1xsolar metallicity

GW150914





GW150914

observed by	LIGO L1, H1	duration from 30 Hz ~ 200 ms		
source type	black hole (BH) binary	# cycles from 30 Hz ~10		
date	14 Sept 2015	peak GW strain 1 x 10 ⁻²¹		
time	09:50:45 UTC	peak displacement of		
likely distance	0.75 to 1.9 Gly	interferometers arms ±0.002 fm		
	230 to 570 Mpc	frequency/wavelength 150 Hz, 2000 km		
redshift	0.054 to 0.136	at peak GW strain		
signal-to-noise ratio	24	peak speed of BHs ~ 0.6 c		
false alarm prob.	less than 1 in 5 million	peak GW luminosity 3.6 x 10 ⁵⁶ erg s ⁻¹		
		radiated GW energy 2.5-3.5 M⊙		
false alarm rate	1 in 200,000 yr	remnant ringdown freq. ~ 250 Hz		
Source Masses M⊙		,		
total mass	65	remnant damping time ~ 4 ms		
chirpmass	28	remnant size, area 180 km, 3.5 x 10 ⁵ km ²		
primary BH	32 to 41	consistent with passes all tests		
secondary BH	25 to 33	general relativity? performed		
remnant BH	62	graviton mass bound $< 1.2 \times 10^{-22} \text{ eV}$		
mass ratio	0.6 to 1	coalescence rate 2 to 400 Gpc ⁻³ yr ⁻¹		
primary BH spin	< 0.7	online trigger latency ~ 3 min		
secondary BH spin	< 0.9	# offline analysis pipelines 5		
remnant BH spin	0.7	~ 50 million (=20 000		
signal arrival time	arrived in L1 7 ms	CPU hours consumed PCs run for 100 days)		
delay	be <mark>f</mark> ore H1			
likely sky position	Southern Hemisphere	papers on Feb 11, 2016 13		
likely orientation resolved to	face-on/off ~600 sq. deg.	# researchers in 15 countries		




Detector noise introduces errors in measurement. Parameters with a range (e.g. distance) are 90% credible bounds; fractional error on parameters without a range is less than 10%. **Acronyms**: L1=LIGO

GW150914: parametr estimation

	EOBNR	IMRPhenom	Overall
Detector-frame total mass $M/{ m M}_{\odot}$	$70.3_{-4.8}^{+5.3}$	$70.7^{+3.8}_{-4.0}$	$70.5^{+4.6\pm0.9}_{-4.5\pm1.0}$
Detector-frame chirp mass $\mathcal{M}/\mathrm{M}_{\odot}$	$30.2_{-1.9}^{+2.5}$	$30.5_{-1.8}^{+1.7}$	$30.3^{+2.1\pm0.4}_{-1.9\pm0.4}$
Detector-frame primary mass $m_1/{ m M}_{\odot}$	$39.4_{-4.9}^{+5.5}$	$38.3_{-3.5}^{+5.5}$	$38.8^{+5.6\pm0.9}_{-4.1\pm0.3}$
Detector-frame secondary mass $m_2/{ m M}_{\odot}$	$30.9_{-4.4}^{+4.8}$	$32.2^{+3.6}_{-5.0}$	$31.6^{+4.2\pm0.1}_{-4.9\pm0.6}$
Detector-frame final mass $M_{\rm f}/{ m M}_{\odot}$	$67.1_{-4.4}^{+4.6}$	$67.4_{-3.6}^{+3.4}$	$67.3^{+4.1\pm0.8}_{-4.0\pm0.9}$
Source-frame total mass $M^{ m source}/{ m M}_{\odot}$	$65.0_{-4.4}^{+5.0}$	$64.6_{-3.5}^{+4.1}$	$64.8^{+4.6\pm1.0}_{-3.9\pm0.5}$
Source-frame chirp mass $\mathcal{M}^{\mathrm{source}}/\mathrm{M}_{\odot}$	$27.9_{-1.8}^{+2.3}$	$27.9_{-1.6}^{+1.8}$	$27.9^{+2.1\pm0.4}_{-1.7\pm0.2}$
Source-frame primary mass $m_1^{ m source}/{ m M}_{\odot}$	$36.3_{-4.5}^{+5.3}$	$35.1_{-3.3}^{+5.2}$	$35.7^{+5.4\pm1.1}_{-3.8\pm0.0}$
Source-frame secondary mass $m_2^{ m source}/{ m M}_{\odot}$	$28.6_{-4.2}^{+4.4}$	$29.5^{+3.3}_{-4.5}$	$29.1^{+3.8\pm0.2}_{-4.4\pm0.5}$
Source-fame final mass $M_{ m f}^{ m source}/{ m M}_{\odot}$	$62.0_{-4.0}^{+4.4}$	$61.6_{-3.1}^{+3.7}$	$61.8^{+4.2\pm0.9}_{-3.5\pm0.4}$
Mass ratio q	$0.79^{+0.18}_{-0.19}$	$0.84^{+0.14}_{-0.21}$	$0.82^{+0.16\pm0.01}_{-0.21\pm0.03}$
Effective inspiral spin parameter $\chi_{\rm eff}$	$-0.09_{-0.17}^{+0.19}$	$-0.03^{+0.14}_{-0.15}$	$-0.06^{+0.17\pm0.01}_{-0.18\pm0.07}$
Dimensionless primary spin magnitude a_1	$0.32^{+0.45}_{-0.28}$	$0.31^{+0.51}_{-0.27}$	$0.31^{+0.48\pm0.04}_{-0.28\pm0.01}$
Dimensionless secondary spin magnitude a_2	$0.57^{+0.40}_{-0.51}$	$0.39^{+0.50}_{-0.34}$	$0.46^{+0.48\pm0.07}_{-0.42\pm0.01}$
Final spin $a_{ m f}$	$0.67^{+0.06}_{-0.08}$	$0.67^{+0.05}_{-0.05}$	$0.67^{+0.05}_{-0.07}^{\pm0.00}_{\pm0.03}$
Luminosity distance $D_{\rm L}/{ m Mpc}$	390^{+170}_{-180}	440^{+140}_{-180}	$410^{+160\pm20}_{-180\pm40}$
Source redshift z	$0.083^{+0.033}_{-0.036}$	$0.093^{+0.028}_{-0.036}$	$0.088^{+0.031\pm0.004}_{-0.038\pm0.009}$
Upper bound on primary spin magnitude a_1	0.65	0.71	0.69 ± 0.05
Upper bound on secondary spin magnitude a_2	0.93	0.81	0.88 ± 0.10
Lower bound on mass ratio q	0.64	0.67	0.65 ± 0.03
Log Bayes factor $\ln \mathcal{B}_{\mathrm{s/n}}$	288.7 ± 0.2	290.1 ± 0.2	_








BBH Formation from Isolated Binaries Spin Tilts

VK 2000



even smaller tilts for more massive BBH and smaller BH kicks

anti-alignment highly unlikely

see also: Gerosa, ..., O'Shaughnessy, Sperhake 2013

On that ordinary Monday ...



On September 14th ... a message from the online GW-burst pipeline

GraceDB Processor

~ 6am ET (

September 14, 2015 5:54 AM

Details

Inbox - UF exchange

To: klimenko@phys.ufl.edu, reed.essick@ligo.org, Marco Drago action required for GraceDB event: G184098 (burst_cwb_allsky)

action required for GraceDB event : https://gracedb.ligo.org//events/view/G184098 (burst_cwb_allsky) cwb eventcreation



On September 14th ...

~7am: "a very interesting event in the last hour"



On September 14th ...

~7am: "a very interesting event in the last hour"

+20min: "no scheduled hardware injection"



~7am: "a very interesting event in the last hour"

+20min: "no scheduled hardware injection"

+10min: "very interesting indeed! looks like a high-mass inspiral?"



~7am: "a very interesting event in the last hour"

+20min: "no scheduled hardware injection"

+10min: "very interesting indeed! looks like a high-mass inspiral?"

+25min: "The Omega scans have finished and I do not see any DQ issues at the time of the trigger. The data looks quite clean at both detectors"



~7am: "a very interesting event in the last hour"

+20min: "no scheduled hardware injection"

+10min: "very interesting indeed! looks like a high-mass inspiral?"

+25min: "The Omega scans have finished and I do not see any DQ issues at the time of the trigger. The data looks quite clean at both detectors"

+15min: "This is clean and very significant inspiral with Mchirp = 27 +- 2 Mo... FAR << 1.e-10 Hz.

If this is not injection,

I guess, we need to do the detection checklist..."



~7am: "a very interesting event in the last hour"

+20min: "no scheduled hardware injection"

+10min: "very interesting indeed! looks like a high-mass inspiral?"

+25min: "The Omega scans have finished and I do not see any DQ issues at the time of the trigger. The data looks quite clean at both detectors"

and off we went on an unexpected ride

+15min: "This is clean and very significant inspiral with Mchirp = 27 +- 2 Mo... FAR << 1.e-10 Hz.

If this is not injection,

I guess, we need to do the detection checklist..."





Text message from my past grad student Ben Farr



Ben

Details

Mon, Sep 14, 5:42 PM

Have you been keeping up on LSC emails today?

Not yet - what should I catch?

About the grb or the loud trigger in ER8?

G18?


Loud trigger baby! This could be the one. I was on most of the telecons this morning if you want an update

Yes :-)

Later on skype?




GW150914






GW150914






GW150914





Advanced LIGO





GW150914: Binary BH merger rate

assumed to be constant within current sensitive volume out to z~0.5

for GW150914-like BBH mergers: 2 - 53 per Gpc^3 per yr

BUT ... search reveals many more triggers (<2 sigma)


what if we assume they are all from the same population accounting for their probability of being astrophysical?

6-400 Gyr^3 per yr




BBH Formation from Isolated Binaries

Belczynski, ..., Bulik, O'Shaughnessy, ..., Holz 2010 Dominik, Belczynski, ..., Holz, ..., Bulik, Mandel, O'Shaughnessy 2012 Dominik, Belczynski, ..., Holz, Bulik, Mandel, O'Shaughnessy 2013



BBH Formation from Isolated Binaries

Belczynski, ..., Bulik, O'Shaughnessy, ..., Holz 2010 Dominik, Belczynski, ..., Holz, ..., Bulik, Mandel, O'Shaughnessy 2012 Dominik, Belczynski, ..., Holz, Bulik, Mandel, O'Shaughnessy 2013



Branching Ratios for this Evolutionary Channel:

99% of BBH at solar metallicity 90% of BBH at 0.1xsolar metallicity


Most important factors:
Stellar Winds & Metallicity
Common-Envelope Treatment
BH Natal Kicks

Other model factors:
Initial binary properties
Stellar models & properties
Mass Transfer
Tides in binaries

...


BBH Formation from Isolated Binaries with SFH & IMR waveforms

Dominik, ..., O'Shaughnessy, Mandel, Belczynski, ..., Holz, Bulik, Pannarale 2015



chirp mass

detectable BBH



mass ratio

BBH from clusters key model elements

All from binary evolution



Cluster Models

Realistic N particles Few-body dynamics/interactions

Most important factors:
Stellar Winds & Metallicity
Common-Envelope Treatment
BH Natal Kicks

Other model factors:


Initial binary properties
Stellar models & properties
Mass Transfer
Tides in binaries

Cluster:
Total Mass
Size
Central Density
Metallicity
Age

Mass Function Volume Density of Clusters


BBH Rates: Field & Clusters

Ziosi, ..., Branchesi, & Tormen 2014



Ready for the Era of Gravitational-Wave Astrophysics?


Stochastic Background from BBH mergers ...





Ready for the Era of Gravitational-Wave Astrophysics?

Stochastic Background from BBH mergers ...




within reach at design sensitivity



Hulse-Taylor Binary Pulsar: The first relativistic binary pulsar

Hulse-Taylor Binary Pulsar: The first relativistic binary pulsar

A radio pulsar with a NS as its binary companion



Hulse-Taylor Binary Pulsar: The first relativistic binary pulsar

A radio pulsar with a NS as its binary companion


PSR 1913 + 16 -- Timing of pulsars

17 / sec

17 / sec

Neutron Binary System
• separated by 10^6 miles
• $m_1 = 1.44m_{\odot}$; $m_2 = 1.39m_{\odot}$; $\epsilon = 0.617$ Prediction from general relativity
• spiral in by 3 mm/orbit
• rate of change orbital period

Rate of orbital decay consistent with gravitational wave losses within 0.3%



Hulse-Taylor Binary Pulsar: The first relativistic binary pulsar

A radio pulsar with a NS as its binary companion

PSR 1913 + 16 -- Timing of pulsars


17 / sec

17 / sec

Neutron Binary System
• separated by 10⁶ miles
• m₁ = 1.44m_Θ; m₂ = 1.39m_Θ; ε = 0.617

Prediction from general relativity
• spiral in by 3 mm/orbit
• rate of change orbital period

Rate of orbital decay consistent with gravitational wave losses within 0.3%



5 more relativistic NS-NS have been discovered

Einstein's gravity: mass warps space



Einstein's gravity: mass warps space



credit: C. Rodriguez

Einstein's gravity: mass warps

space

warped space tells mass how to move





credit: C. Rodriguez

Einstein's gravity: moving warps

generate waves!







credit: C. Rodriguez

Einstein's gravity: moving warps

generate waves!








Initial LIGO: Binary Coalescence Upper Limits

Abadie et al 2012




BH mass: 5 Msun 90% Upper Limits



Initial LIGO: Binary Coalescence Upper Limits


Abadie et al 2012




BH mass: 5 Msun 90% Upper Limits


BH mass: 20 Msun







$$\frac{\Delta x}{x} = \frac{1}{2} h_{xx}$$







$$\frac{\Delta x}{x} = \frac{1}{2} h_{xx}$$







$$\frac{\Delta x}{x} = \frac{1}{2} h_{xx}$$








$$\frac{\Delta x}{x} = \frac{1}{2} h_{xx}$$





very weak:

$$h \sim \frac{1}{r} \frac{G}{c^4} \frac{MR^2}{T^2} \sim \frac{GM/c^2}{r} \frac{u^2}{c^2} \sim \frac{R_{sch}}{r} \left(\frac{u}{c}\right)^2$$

ideally: massive, highly compact, fast moving, nearby objects

very weak:

$$h \sim \frac{1}{r} \frac{G}{c^4} \frac{MR^2}{T^2} \sim \frac{GM/c^2}{r} \frac{u^2}{c^2} \sim \frac{R_{sch}}{r} \left(\frac{u}{c}\right)^2$$

ideally: massive, highly compact, fast moving, nearby objects

direct detection goal:

measure GW strain -> relative distance changes

very weak:

$$h \sim \frac{1}{r} \frac{G}{c^4} \frac{MR^2}{T^2} \sim \frac{GM/c^2}{r} \frac{u^2}{c^2} \sim \frac{R_{sch}}{r} \left(\frac{u}{c}\right)^2$$


ideally: massive, highly compact, fast moving, nearby objects

direct detection goal:

measure GW strain -> relative distance changes


Black Hole of 10 solar masses at the speed of light located at the Galactic center: $h \sim 10^{-17}$ at the Virgo cluster: $h \sim 10^{-20}$

GW150914: first-ever strong-field GR tests




GW150914: strong-field GR tests


after NR waveform is removed, residuals are consistent with noise



GW150914: strong-field GR tests

final-BH properties consistent with and without including strong-field GR models







GW150914: strong-field GR tests

First ever high PN-order limits on deviations from GR

fractional deviation from GR 10^3 10^2 10^{1} $\delta\hat{arphi}$ 10^{0} 10^{-1} GW150914 10^{-2} J0737-3039 0PN 0.5PN 1PN 1.5PN 2PN 2.5PN 3PN 3.5PN PN order

 $1PN \sim (v/c)^2$



Template Analysis: measuring deviations

Variable G

Scalar Field

$$\Psi(f) = 2\pi f t_c - \phi_c - \frac{\pi}{4} + \frac{3}{128} u^{-5} \left[1 - \frac{25}{1536} \dot{G} \mathcal{M} u^{-8} - \frac{5}{84} \frac{S^2}{\omega_{BD}} \eta^{3/5} u^{-2} \right]$$

$$+ \left(\frac{3715}{756} + \frac{55}{9} \eta \right) \eta^{-2/5} u^2 - 16\pi \eta^{-3/5} u^3 - \frac{128}{3} \frac{\pi^2 D \mathcal{M}}{\lambda_q^2 (1+z)} u^2 + \dots \right]$$

Massive Graviton

$$u = (\pi \mathcal{M}f)^{1/3}$$

$$\Psi(f) = \Psi_{GR}(f) + \sum_{i} \beta_i u_i^b$$