

Professor, Dept. of Earth and Environmental Sciences


Wesleyan University, Middletown CT


wesleyan.edu/planetary


Planetary Science Decadal Survey: Venus Panel March 24, 2021


Credit: NASA GSFC

### Tesserae and Mountains











#### GENERALIZED EMISSIVITY CLASSIFICATION FOR ALL VENUS HIGHLANDS



# Interpretation 1: the Tesserae are granitoids formed the common way

- Most Earth granitoids formed via melting of continental crust, sediments and/or peridotitic mantle in the presence of H<sub>2</sub>O [e.g., Campbell and Taylor, 1983; Brown, 2013]. Settings:
  - · Fractionation of hydrous mantle above subducting plate
  - Remelt older continental crust or seds by basaltic underplating or collision (e.g., Hubbert and Sparks, 1988, Thompson and Connolly, 1985)
  - Archean TTGs partial melting of hydrated basalts/eclogites (e.g., Martin et al., 2005)
  - Rare (~10<sup>4</sup> km<sup>3</sup>) felsic eruptions on Earth associated with continental breakup (plumes) - such volumes requires the partial melting of hydrous lower continental crust materials [e.g., Bryan et al., 2002].
- To get <u>continents' worth</u> of granitic magma need **water** and **recycling**. Recycling invoked in Venus's past (Parmentier and Hess, 1992; Turcotte et al, 1999) and present (Elkins-Tanton et al., 2007). Therefore, <u>likely limited to lifetime of water on Venus Gilmore PSDS</u>



Felsic Rocks:
Need water and lithosphere cycling

## Interpretation 2: the Tesserae are granitoids formed via differentiation of large mafic bodies


- Process should generate ~10% felsic melt (e.g., Spulber and Rutherford, 1983; Shelnutt, 2013)
- LIPs typically associated with ~5% felsic rocks (Bryan et al. 2002).

#### SHELLNUTT: FORMATION OF SILICIC ROCKS ON VENUS

**Table 2.** Physical Parameters of the Modeled Silicic Liquids<sup>a</sup>


Sample	Process	Pressure (GPa)	Temperature (°C)	Viscosity (Pa s)	Density (g/cm <sup>3</sup> )	Rock Type	Water State
Venera 13-1	FC	0.01	1060	10 <sup>3.8</sup>	2.44	Phonolite	anhydrous
Venera 13-2	FC	0.1	1070	$10^{3.7}$	2.46	Phonolite	anhydrous
Venera 13-3	FC	0.01	1060	$10^{3.3}$	2.43	Phonolite	hydrous
Venera 13-4	FC	0.1	1060	$10^{3.5}$	2.44	Phonolite	hydrous
Venera 14-1	FC	0.01	970	$10^{5.8}$	2.42	Rhyolite	anhydrous
Venera 14-2	FC	0.1	980	$10^{5.5}$	2.44	Rhyolite	anhydrous
Venera 14-3	FC	0.01	960	$10^{5.0}$	2.37	Rhyolite	hydrous
Venera 14-4	FC	0.1	910	$10^{4.8}$	2.32	Rhyolite	hydrous
Venera 14-5	FC	1	900	$10^{3.8}$	2.49	Rhyolite	hydrous
Venera 14-6	PM	0.01	1080	$10^{2.6}$	2.54	Andesite	hydrous
Venera 14-7	PM	0.1	940	$10^{3.3}$	2.27	Andesite	hydrous
Venera 14-8	PM	1	880	$10^{4.6}$	2.49	Rhyolite	hydrous

<sup>&</sup>lt;sup>a</sup>FC = fractional crystallization, PM = partial melting.



3/24/21

### Interpretation 3: the Tesserae are mafic.....but weathered differently



Elevate pH<sub>2</sub>O get everything to completion and more Weathering during plains emplacement? On ancient Venus?


### Interpretation 4: the Tesserae are not igneous





3/24/21


### **Morphological Evidence for Layering**

Byrne et al., 2020





















Khawja et al., 2020



### **Constraints from Buckling Theory**

- · Tesserae deformed in a different thermal regime
- Characteristic wavelengths of structures are similar across tesserae and consistent with a lithosphere that is thinner than that estimated today (Brown and Grimm, 1997; 1999).
- Strain rate estimates 1-2 orders of magnitude > today (Grimm, 1994).
- · A global process.

3/24/21



Venus present Venus tessera 50 70 10 b. Columbia diabase (model, 48% SiO<sub>2</sub>) -15 -30 -60 c. Venera 14 (model, 49% SiO<sub>2</sub>) -15 -30 0 0.5 1 10 30 50 70 d. quartz monzonite (model, 60%  ${
m SiO}_2$ , 10% quartz) -14 -15 -30 -60 0 0.5 1 10 30 5 e. granite (model, 72% SiO<sub>2</sub>, 31% quartz) log(10% strain time) depth (km) log(strain re 0.5 30 50 geotherm (K/km)  $\sigma_1$ -  $\sigma_2$  (GPa)

a. Columbia diabase (experimental, 48% SiO<sub>2</sub>)

-30

-15

Brown and Grimm, 1999

Gilmore - PSDS

Resor, Gilmore et al., JGR, in press

# Tesserae are key to understanding Venus geologic evolution, climate and habitability

I encourage the Decadal to Emphasize and Prioritize:

- Tessera Composition
  - Mineralogy from orbit and probe VERITAS/DAVINCI+
  - Measure chemistry and mineralogy at tessera landing site VITaL/Flagship
  - Experimental work on weathering of rocks under Venus conditions
- Tessera Morphology and History
  - High resolution imaging and topography (CTX, MOC, MOLA at Venus) VERITAS/DAVINCI+
  - Improved gravity VERITAS

A Venus Flagship class mission would enable direct measurement of tessera terrain and a comprehensive study of the inventory, movement and fate of the volatiles in the current and ancient Venus system. I urge the Decadal to recommend a Venus Program of missions, research and technology advancement, including a Flagship-class mission in the next decade.