



Distribution of Awarded Observing Time



2018B ~30% of observing time to small-body science 2020B



Telescope Operations

- National facility with observing time open to all through peer review of proposals
- 50% of observing time dedicated to solar system observations and mission support
- Flexible scheduling (multiple programs per night), 90% of observing is now remote
- Target-of-opportunity (ToO) programs
- Director Discretionary Time and NASA programmatic requests
- Daytime observing as necessary
- Dedicated time available for observing campaigns (comets, mission events)

Small-Body Mission Support

- Observations for mission planning and support (imaging and spectroscopy)
 - Characterization and support observation of mission targets

Eros (NEAR)
Vesta and Ceres (Dawn)

- Jupiter Trojans (Lucy) - Psyche (Psyche)

- Itokawa (Hayabusa) - Ryugu (Hayabusa 2)

- Bennu (OSIRIS-REX) - Phaethon / Geminid meteor shower (DESTINY+)


Didymos (DART)
possible PHO mission targets

- Stellar occultations by dwarf planets (context for New Horizons)

Seasonal changes on Pluto (New Horizons)

Mission events


- Deep Impact: impact on comet 9P/Tempel 1
- LCROSS: search for volatiles released on lunar impact




Critical capabilities for NEO characterization

- ToO rapid response for newly discovered targets
- Accurate non-sidereal tracking at rates up to 60"/sec (1°/min)
- Multi-instrument characterization (SpeX + MORIS + MIRSI +...)
- · Adding wide-field finder to improve acquisition and tracking







Data courtesy Nick Moskovitz

NEO Science

Compositional Distributions and Evolutionary Processes for the Near-Earth Object Population: Results from the MIT-Hawaii Near-Earth Object Spectroscopic Survey (MITHNEOS)


R. P. Binzel + 20 co-authors (*Icarus* 2019)

Study includes:


- ~1000 NEOs, database http://smass.mit.edu/smass.html
- ~350 Mars-crossing asteroids
- Large portion of the data were obtained with SpeX (prism mode, 0.8 – 2.5 um, taken 2000 – present)

Topics addressed in analysis:

- Phase reddening
- Taxonomic distributions
- Space weathering trends
- Shock darkening trends
- Ordinary chondrite mineralogy
- NEO / meteorite source regions



Distribution of targets included in MITHNEOS study. The three panels depict different distance scales with markers representing target diameters (marker size), albedo (color), and geocentric position at the time of observation.




Trot = 134 K

3.00 μm-

3350

Kuiper Belt



Planned Future Capabilities for Small-Body Science (2023-2032)

- MIRSI upgrade for simultaneous mid-IR and visible photometry albedos and diameters (2021)
- Wide-angle telescope finder to acquire NEOs with large position uncertainties (2021)
- Low-order off-axis WFS to improve image quality (2022)
- Proposal for new facility instrument optimized for small-body science (SPECTRE)

MIRSI Upgrade

5-20 µm camera and spectrograph upgrade

Replace LN2/LHe with closed-cycle cooler

New low-noise array controller

New hexapod chopper (no top-end change)

Add CCD channel for simultaneous N+V

 NEO albedos and diameters: ~200 per year with 10 hours observing per month





Wide-angle Finder ('Opihi)

Capabilities

- 0.43-m telescope
- 0.5 degree FOV (2k² Andor CCD)
- V=19.5 60s-20*σ*
- Filters *g*, *r*, *i*, *z*
- Boot-strap photometry for all instruments
- Extinction and cloud monitor
- Context camera
- On-telescope for testing in fall 2021



SPECTRE: 0.4-4 µm R=150 IFU SPECTROGRAPH

Endorsed by IRTF community workshop (2018)

Enacted by Intil Community Workshop (2010)


Optical, NIR and MIR channels

Technical features

- 7" x 7" IFU absolute spectro-photometry on point sources
- High throughput (70%), no slit losses
- Compact and rigid spectrograph, no moving optics
- 3'x3' CCD camera for A&G (e.g. NEOs)

Optimized for small-body science


- R=150 for solid-state features and SEDs
- One-shot 0.4-4 µm to remove degeneracy, time variability
- > 1 mag improvement in sensitivity, multiplex advantage
- Rapid response for newly discovered NEO targets
- Many more targets observable throughout Solar System
- Seeking funding from NSF or NASA (\$3.5m)



2.0

1.0

Data courtesy of Driss Takir



3.5

3.0



SPECTRE Mission Support for Lucy Trojan Asteroid Tour (2027-2033)



3548 Eurybates (flyby 8/27)

- 64 km diameter, rotation 9 hours
- 0.4-4 μ m R=100 50 σ in 1 hour at opposition
- map at sub-hemispheric resolutions
- 50 Trojans brighter than Eurybates
- order of magnitude more objects observable 0.4-2.4 μm

Ground-truth Lucy observations can then be applied to a large population of Trojans

