
# Creating policy-relevant evidence for population health science: promise and pitfalls


Sandro Galea



It all started here

# Cholesterol distributions and coronary heart disease rates, men 30–59 years of age, North Karelia, Finland, 1972–1992





Emerging experiments in population health



### **Press Releases**

Boston Medical Center to Invest \$6.5 Million in Affordable Housing to Improve Community Health and Patient Outcomes, Reduce Medical Costs

December 07, 2017

For More Information, Contact:
David Kibbe
Office of Communications
617-638-8499
david.kibbe@bmc.org

## Boston Medical Center to Invest \$6.5 Million in Affordable Housing to Improve Community Health and Patient Outcomes, Reduce Medical Costs

BMC's investment, the first in Boston, joins a growing national trend of hospitals prescribing housing for health

(Boston) – Dec. 7, 2017 – Boston Medical Center is investing \$6.5 million over five years to support a wide range of affordable housing initiatives, in an innovation lab approach that will be studied closely to determine the best ways that health care systems can improve both community and patient level health and reduce medical costs by addressing homelessness and housing

We do have some evidence

Q

CDC A-Z INDEX Y

### Office of the Associate Director for Policy

### AD for Policy Home

Health System Transformation

6|18 Initiative

Health Impact in 5 Years

#### The HI-5 Interventions

HI-5 14 Evidence-Based Community-Wide Interventions Slides

HI-5 Stories from Public Health Innovators

School-Based Programs to Increase Physical Activity

School-Based Violence Prevention

Safe Routes to School (SRTS)

Motorcycle Injury Prevention

Tobacco Control Interventions

Access to Clean Syringes

**Pricing Strategies for** Alcohol Products

Multi-component Worksite Obesity Prevention

Early Childhood Education

Clean Diesel Bus Fleets

**Public Transportation** System Introduction or <u>AD for Policy Home</u> > <u>Health System Transformation</u> > <u>Health Impact in 5 Years</u>

### The HI-5 Interventions







### What are the HI-5 Interventions?



### HEALTH IMPACT IN 5 YEARS

### Interventions Changing the Context


- School-Based Programs to Increase Physical Activity
- School-Based Violence Prevention
- Safe Routes to School
- · Motorcycle Injury Prevention
- Tobacco Control Interventions
- Access to Clean Syringes
- Pricing Strategies for Alcohol Products
- Multi-Component Worksite Obesity Prevention

### Interventions Addressing the Social Determinants of Health

- Early Childhood Education
- Clean Diesel Bus Fleets
- Public Transportation: System Introduction or Expansion
- Home Improvement Loans and Grants
- Earned Income Tax Credits
- Water Fluoridation







acs, J.B., Cost officering Investments in Children Isaacs. 2007, The Brookings Institute: Washington D.C.



# LEVERAGING THE SOCIAL DETERMINANTS OF HEALTH: WHAT WORKS?

## **Executive Summary**

**JUNE 2015** 




prepared for the Blue Cross Blue Shield of Massachusetts Foundation by Lauren A. Taylor, Caitlin E. Coyle, Chima Ndumele, Erika Rogan, Maureen Canavan, Leslie Curry, and Elizabeth H. Bradley

Challenges in producing evidence

1. External validity


Let us suppose we want to know: Does a particular intervention result in normotension?

### Sample 1 from Population 1:




Risk difference 
$$=\frac{50}{100} - \frac{200}{400} = 0.5 - 0.5 = 0 (95\% \text{ CI} - 0.109, 0.109)$$

### Sample 1 from Population 1:




### Sample 2 from Population 2:




Risk difference 
$$=\frac{100}{200} - \frac{50}{300} = 0.5 - 0.166 = 0.333 (95\% CI 0.252, 0.414)$$

### Sample 2 from Population 2:




### Sample 3 from Population 3:



Risk difference 
$$=$$
  $\frac{40}{150} - \frac{60}{350} = 0.266 - 0.171 = 0.095 (95\% CI 0.014, 0.176)$ 

### Sample 3 from Population 3:



### Why?


The intervention alone does not cause normotension; the intervention needs to happen together with exposure to other conditions

What the causal structure may look like if X, the intervention, and Y, social condition are both necessary and insufficient causes of disease

Component Causes		
X	Υ	
<b>✓</b>	<b>✓</b>	
<b>✓</b>		
	<b>✓</b>	

What the causal structure may look like if X, the intervention, and Y, social condition are both necessary and insufficient causes of disease


Component Causes		Probability of disease
X	Υ	
<b>✓</b>	•	1
<b>✓</b>		0
	<b>~</b>	0
		0



Unexposed


Sample 1 from Population 1:

200




200

Sample 2 from Population 2:



Risk difference  $=\frac{100}{200} - \frac{50}{300} = 0.5 - 0.166 = 0.333 (95\% CI 0.252, 0.414)$ 


Sample 3 from Population 3:



Risk difference =  $\frac{40}{150} - \frac{60}{350} = 0.266 - 0.171 = 0.095 (95\% CI 0.014, 0.176)$ 

0% social condition

Sample 1 from Population 1:




Risk difference  $=\frac{50}{100} - \frac{200}{400} = 0.5 - 0.5 = 0 \text{ (95\% CI} - 0.109, 0.109)$ 

50% social

condition


Sample 2 from Population 2:



 $\mbox{Risk difference} \ = \frac{100}{200} - \frac{50}{300} = 0.5 - \ 0.166 \ = \ 0.333 \ (95\% \ \mbox{CI} \ 0.252, 0.414)$ 

40% social condition

Sample 3 from Population 3:





Risk difference =  $\frac{40}{150} - \frac{60}{350} = 0.266 - 0.171 = 0.095 (95\% CI 0.014, 0.176)$ 


0% social condition

50% social condition











No one here will develop normotension due to the intervention because the causal partner is not present; hence equal likelihood of normotension among those exposed/unexposed to the intervention

	onent uses	Probability of diseas
Χ	Υ	
<b>✓</b>	<b>✓</b>	1
<b>V</b>		0
	~	0
		0





50% of those who are exposed to the intervention exposed to the social condition, hence have normotension

Sample 2 from Population 2:


Diseased Not Diseased

Exposed 100 100


Unexposed 50 250

Risk difference  $=\frac{100}{200} - \frac{50}{300} = 0.5 - 0.166 = 0.333 (95\% CI 0.252, 0.414)$ 


40% of those who are exposed to the intervention are exposed to the social condition, hence have normotension



17% of those who are unexposed to the intervention have normotension




17% of those who are unexposed to the intervention have normotension



Therefore under a very plausible assumption of cooccurring causes, the medical intervention-health association can only be understood if we understand the other factors that distinguish between samples

## 2. Ubiquity

Figure 1. A metaphor for ubiquity



The goldfish are surrounded by water and everything they do is influenced by the quality of the water in which they live; therefore, water is a ubiquitous factor influencing the fish and needs to be taken into consideration every time we may want to improve the lives of the fish.

## Crack Babies: The Worst Threat Is Mom Herself

September 17, 1989

Crack's Toll Among Babies: A Joyless View

September 6, 1988

**Cocaine: Litany of Fetal Risks Grows** 

By Douglas J. Besharov


AST WEEK in this city, Greater Southeast Community Hospital released a 7-week-old baby to her homeless, drug-addicted mother even though the child was at severe risk of pulmonary arrest. The hospital's explanation: "Because [the mother] demanded that the baby be released."

The hospital provided the mother with an apnea monitor to warn her if the baby stopped breathing while asleep, and trained her in CPR. But on the very first night, the mother went out drinking and left the child at a friend's house-without the monitor. Within seven hours, the baby was dead. Like Dooney Waters, the 6year-old living in his mother's drug den, whose shocking story was reported in The Washington Post last

week, this child was all but abandoned by ities.



CHILDREN OF COCAINE (By Charles Krauthammer)



Predictor for Peabody Picture Vocabulary Test score	Coefficient	P-value	
Gestational cocaine exposure	-2.89	0.26	
Assessment no.	2.72	<0.001	
Gestational cocaine exposure x assessment no.	0.58	0.51	
Age at 1st assessment	-0.36	0.76	
Female gender	-4.93	0.058	
Parental nurturance	-0.31	0.89	
Environmental stimulation	5.91	0.039	
Caregiver BDI-II depression score	0.03	0.84	


How much of our cognitive ability is determined by our genes?






**=** GE+











































Is this all theoretical?

Proportion of total Full-Scale IQ variance accounted for by A, C, and E plotted as a function of observed socioeconomic status (SES).



3. We cannot think in dichotomies




Figure 3 Percentage distribution of serum cholesterol levels (mg/dl) in men aged 50–62 who did or did not subsequently develop coronary heart disease (Framingham Study<sup>5</sup>)







Figure 3 Percentage distribution of serum cholesterol levels (mg/dl) in men aged 50–62 who did or did not subsequently develop coronary heart disease (Framingham Study<sup>5</sup>)




**FIGURE 2.** Probability distributions of a marker, X, in cases (solid curves) and controls (dashed curves) consistent with the logistic model logit $P(D=1|X)=\alpha+\beta X$ . It has been assumed that X has a mean of 0 and a standard deviation of 0.5 in controls so that a unit increase represents the difference between the 84th and 16th percentiles of X in controls. The marker is normally distributed, with the same variance in cases. The odds ratio (OR) per unit increase in X is shown.

Pepe MS, Janes H, Longton G, Leisenring W, Newcomb P. Limitations of the odds ratio in gauging the performance of a diagnostic, prognostic, or screening marker. *American Journal of Epidemiology* 2004; 159:882-890.







Meigs JB, Shrader P, Sullivan LM, McAteer JB, Fox CS, Dupuis J, Manning AK, Florez JC, Wilson PW, D'Agostino RB Sr, Cupples LA. Genotype score in addition to common risk factors for prediction of type 2 diabetes. N Engl J Med. 2008 Nov 20;359(21):2208-19




Meigs JB, Shrader P, Sullivan LM, McAteer JB, Fox CS, Dupuis J, Manning AK, Florez JC, Wilson PW, D'Agostino RB Sr, Cupples LA. Genotype score in addition to common risk factors for prediction of type 2 diabetes. N Engl J Med. 2008 Nov 20;359(21):2208-19




**Figure 1** Risk ratio for the effect of gene on disease across prevalences of environmental variables and background rate of disease. G=genetic cause of disease. E=environmental cause of disease; the environmental cause requires presence in order for the genetic marker to have an effect. X=background rate of the disease—all causes that are not either G or E. Graph: y axis=risk ratio for the effect of genetic marker on disease; x axis=prevalence of genetic marker in each population.



**Figure 1** Risk ratio for the effect of gene on disease across prevalences of environmental variables and background rate of disease. G=genetic cause of disease. E=environmental cause of disease; the environmental cause requires presence in order for the genetic marker to have an effect. X=background rate of the disease—all causes that are not either G or E. Graph: y axis=risk ratio for the effect of genetic marker on disease; x axis=prevalence of genetic marker in each population.



**Figure 1** Risk ratio for the effect of gene on disease across prevalences of environmental variables and background rate of disease. G=genetic cause of disease. E=environmental cause of disease; the environmental cause requires presence in order for the genetic marker to have an effect. X=background rate of the disease—all causes that are not either G or E. Graph: y axis=risk ratio for the effect of genetic marker on disease; x axis=prevalence of genetic marker in each population.




**Figure 1** Risk ratio for the effect of gene on disease across prevalences of environmental variables and background rate of disease. G=genetic cause of disease. E=environmental cause of disease; the environmental cause requires presence in order for the genetic marker to have an effect. X=background rate of the disease—all causes that are not either G or E. Graph: y axis=risk ratio for the effect of genetic marker on disease; x axis=prevalence of genetic marker in each population.

Grounding this in a conceptual framework

- 1. Population health manifests as a continuum.
- 2. The causes of differences in health across populations are not necessarily an aggregate of the causes of differences in health within populations.
- 3. Large benefits to population health may not improve the lives of all individuals.
- 4. The causes of population health are multilevel, accumulate throughout the life course, and are embedded in dynamic interpersonal relationships.
- 5. Small changes in ubiquitous causes may result in more substantial change in the health of populations than larger changes in rarer causes.
- 6. The magnitude of an effect of exposure on disease is dependent on the prevalence of the factors that interact with that exposure.
- 7. Prevention of disease often yields a greater return on investment than curing disease after it has started.
- 8. Efforts to improve overall population health may be a disadvantage to some groups; whether equity or efficiency is preferable is a matter of values.
- We can predict health in populations with much more certainty than we can predict health in individuals.


The past as prologue

# Simple approaches, a foundational myth





## The effectiveness of simple approaches?



(668 Deaths: Onset date unknown = 127)

twitter/@sandrogalea

sgalea@bu.edu