


A data-driven approach for personalized dietary interventions

Tal Korem (@tkorem)

Departments of Systems Biology & OB/GYN

Columbia University





Prof. Eran Segal





Dr. David Zeevi



# Studying healthy nutrition through postprandial glycemic responses

# Postprandial (post-meal) glucose response as a measure of healthy nutrition





# Glucose responses provide an immediate high-resolution feedback to every meal





Weight	Post-meal blood glucose levels
Risk factor for many diseases	Risk factor for many diseases
Indirect measure of intervention	Direct measurement of meal effect
Periodic	Immediate
Low-resolution	High-resolution
Few data points per participant	Multiple data point per participant on short time-scales


Maintaining normal blood glucose levels is key to fighting metabolic disease...

What's the best way to do this?

### **The Personalized Nutrition Project: Cohort statistics**

3%

8



# Collection of glycemic, nutritional, clinical and microbiome data on a large healthy cohort

### **Initiation**

Gut microbiome 16S rRNA Metagenomics







Using a subcutaneous sensor (iPro2)

130K hours, 1.56M glucose measurements







**Diary** (food, sleep, physical activity) Using smartphone-adjusted website

5,435 days, 46,898 meals, 9.8M Calories, 2,532 exercises

#### **Questionnaires**


Food frequency Lifestyle Medical







### **Real-time food logging**



### Benefits of real-time logging

- Consumption matched in real-time to actual products, brands, etc.
- Serving sizes considered in real-time
- "Live" database
- Real-time compliance monitoring
- Generally improved compliance

### Collection of glycemic, nutritional, clinical and microbiome data on a large healthy cohort

### Recruitment

**Gut microbiome** 16S rRNA **Metagenomics** 





Continuous glucose monitoring

Using a subcutaneous sensor (iPro2)

130K hours, 1.56M glucose measurements







**Diary** (food, sleep, physical activity) Using smartphone-adjusted website

5,435 days, 46,898 meals, 9.8M Calories, 2,532 exercises

#### Questionnaires







#### Standardized meals (50g available carbohydrates)

Day 1 Day 2 Day 3 Day 4















**Anthropometrics** 




Bread & Bread & Glucose Glucose Fructose butter butter

Why would anyone go through this?

### What will I learn?

My response to food




Know my microbiome




In the future: Plan a better diet




### Full accessibility to collected data




### Participant analysis: nutrition, activity & blood tests







### Participant analysis: Overview of microbiota composition



### Participant analysis: Glucose spiking foods







### Participants engagement and retention

- Better data
- Easier trial management
- Minimal recruitment effort
  - Finished recruitment of 900 individuals with a wait-list of 3,000


# What is the variability across people in the response to the same food?


# Different people have widely different post-meal responses to the same standardized meal





# Significant personalized response which increases with food complexity





Standard normal
Measured GI

### Results replicated and validated by subsequent studies





Original Investigation | Diabetes and Endocrinology

## Assessment of a Personalized Approach to Predicting Postprandial Glycemic Responses to Food Among Individuals Without Diabetes

Helena Mendes-Soares, PhD; Tali Raveh-Sadka, PhD; Shahar Azulay, BSc; Kim Edens, MSc; Yatir Ben-Shlomo, BSc; Yossi Cohen, MSc; Tal Ofek, PhD; Davidi Bachrach, BSc, MBA; Josh Stevens, BSc; Dorin Colibaseanu, MD; Lihi Segal, LLB, MBA; Purna Kashyap, MBBS; Heidi Nelson, MD

See corresponding editorial on page 955.

### Estimating the reliability of glycemic index values and potential sources of methodological and biological variability<sup>1–3</sup>

Nirupa R Matthan, <sup>4</sup>\* Lynne M Ausman, <sup>4</sup> Huicui Meng, <sup>4</sup> Hocine Tighiouart, <sup>5</sup> and Alice H Lichtenstein <sup>4</sup>

<sup>&</sup>lt;sup>4</sup>Jean Mayer USDA Human Nutrition Research Center on Aging, Tufts University, Boston, MA; and <sup>5</sup>Tufts Clinical and Translational Science Institute, Tufts Medical Center, Boston, MA

### General recommendations in nutrition are suboptimal















Source: USDA

Can we predict the personal post-prandial glucose response to any complex meal?


### **Prediction scheme**



### Accurate predictions of personalized glucose responses




### Replication and validation by subsequent studies




Can glucose response predictions improve post-prandial glucose responses?

# Constructing personally tailored diets that achieve normal post-prandial glucose responses


- 12 subjects, mostly prediabetics
- Profiled for one week
- Self-selected isocaloric meals, categorized as good / bad
- Good and bad diets assigned double-blindedly, one week each




# A 'good' meal for one person can be a 'bad' meal for another



# Personally tailored diets reduce the post-prandial glucose response



### Personally tailored diets improve post-meal responses




Is this	useful	for l	ong-te	rm in	terver	ntions?


### Challenges with translation to interventions

- Scores a meal rather than provide insights
  - Really not what people expect
  - Not straight forward to devise interventions
  - Accounting for preferences is challenging
  - Maintaining caloric target is external to prediction framework
- Strong assumptions on causality
  - Probably easier to make than for "long-term" outcomes
- Requires extensive variables
  - Real potential of using the gut microbiome

### Long term, algorithm based intervention




# Long term "algorithm-diet" achieves greater glycemic normalization




### Rich data-sets facilitates multiple datadriven investigations


### Predicting caffeine levels from diet data




## Our predictive models explain over 10% of the variance for 543 metabolites




# Metabolites associated with sourdough consumption increase after sourdough (but not white bread) intervention



# Metabolites associated with sourdough consumption increase after sourdough (but not white bread) intervention




### **Summary**



- High interpersonal variability in post-meal glucose response to identical meals
- Personal and microbiome features enables accurate glucose response prediction
- Personalized dietary interventions
   successfully lower post-meal glucose

### **Summary**



- Importance of real-time logging
- Benefits of participant engagement and data-sharing
- The challenges of using this approach for long-term dietary interventions

### **Acknowledgements**

Prof. Eran Segal

Dr. David Zeevi

Dr. David Israeli

**Noam Bar Adina Weinberger** 

Prof. Eran Elinav

**Kfar Shaul Medical Center** 

Omer Weissbrod

Prof. Zamir Halpern

Noam Bar

**Niv Zmora** 

**Tel-Aviv Sourasky Medical Center** 

Orly Ben-Yacov Dar Lador

Iemal Ali Mahdi

Jotham Suez

Tali Avnit-Sagi

Gili 7ilberman

Maya Lotan-Pompan

Lenka Dohnalova

Meirav Pevsner-Fischer

IMI DIRECT consortium

Dr. Tim D. Spector

Flad Matot

Daphna Rothschild

Gal Malka

Noa Kosower

Michal Rein

Rony Bikovsky

Sigal Leviatan

Prof. Avraham Levy

Ben Z. Goldberg







