Allostatic Load and Breast Cancer: Tracing Social Determinant of Health Pathways

Samilia Obeng-Gyasi MD MPH
Endowed Surgical Professor in Health Equity
Associate Professor of Surgery
The Ohio State University
Department of Surgery
Division of Surgical Oncology

The James





Disclosures

No financial disclosures


Funding

American Cancer Society (RSG-22-106-01-CSCT)

Conquer Cancer Breast Cancer Research Foundation Advanced Clinical Research Award (ACRA) for Diversity and Inclusion in Breast Cancer Research

Society of University Surgeons Junior Faculty Award for the Underrepresented in Medicine

Socioenvironmental Factors and Breast Cancer




What is the mechanistic pathway from socioenvironmental exposures to cancer initiation, progression and metastasis?





Ecosocial Theory

Embodiment



Relationship between the body and environment





Agency and Accountability



Acknowledgement of Structural determinants of health as major driver of disease distribution

Weathering Hypothesis


"weathering is the accumulation and culmination of life experiences that are structured by historical and ongoing systemic racism and classism."

-Arline Geronimus PhD

Example: Maternal outcomes between African Immigrants, women racialized as Black and white



Socioenvironmental Stressors and the Stress Pathway





Allostatic load: physiologic dysregulation secondary to exposure to chronic stress.


Allostasis: "maintain stability or homeostasis through change."



Patterns of Allostatic Load



Repeated events of various types



Failure to adapt to the same stressor




Failure to terminate a response



Inadequate release of stress hormones


Measurements of Allostatic Load



Primary mediators (e.g., cortisol)



Secondary outcomes of primary mediators (e.g., systolic blood pressure)



Tertiary outcomes (e.g., hypertension)

Allostatic Load Biomarkers

Primary Mediators

Cortisol


DHEA

Epinephrine


Norepinephrine

Secondary Outcomes		
Cardiovascular	Metabolic	Immune
Systolic blood pressure	A1c	C-reactive protein
Diastolic blood pressure	Albumin	WBC
Heart rate	ВМІ	IL-6
Cholesterol	eGFR	TNF-alpha
HDL	Creatinine	
Triglycerides	Blood glucose	
LDL	Waist-hip-ratio	


Tertiary Outcomes	
Diabetes	
Hypertension	
Obesity	
Cancer	



Allostatic Load Conceptualization



Elevated Allostatic Load and Socioenvironmental Stressors







Allostatic Load and Cancer

Socioenvironmental Stressors and Cancer




- Social isolation
- 2. Neighborhood contextual factors
- 3. Financial instability
- 4. Healthcare system



Hypothalamic-pituitaryadrenal axis + Sympathetic adrenal medullary pathway

Embodiment/Weathering



Immune Dysregulation


↑Immunosuppression

- ↑chronic inflammation
- **↓**Immunoprotection

- 1. Cancer initiation
 - 2. Progression
- 3. treatment tolerability
- 4. Metastasis and mortality

*genetic risks, epigenetic changes


Association of Allostatic Load With All-Cause Mortality in Patients With Breast Cancer



Obeng-Gyasi et al, Association of Allostatic Load With All-cause Mortality Among Patients With Breast Cancer, JAMA Network Open, May 2023

The Implications of Racialized Economic Segregation and Allostatic Load on Mortality in Patients with Breast Cancer

Triple Negative Breast Cancer




18.2% versus 11.6%, p < 0.001

High AL



OR 1.55, 95% CI 1.36-1.76

Worse Mortality



1.41, 95% CI 1.08-1.83


Chen et al, The Implications of Racialized Economic Segregation and Allostatic Load on Mortality in Patients with Breast Cancer, Annals of Surgical Oncology, November 2023

Association Between Neighborhood Opportunity, Allostatic Load, and All-Cause Mortality in Patients With Breast Cancer

Population Exposure Outcomes Environ*AL aHR 1.45, 95% CI 1.11-1.89 Mediation **AL 9.3%** AL +CCI 11.4% Women stage I-III breast cancer aOR 1.21, 95% CI 1.05-1.40 **Neighborhood Opportunity** N = 4089

> Chen et al Association Between Neighborhood Opportunity, Allostatic Load, and All-Cause Mortality in Patients With Breast Cancer, Journal of Clinical Oncology, 2024

Socioenvironmental Stressors and the Stress Pathway







Future Directions










AL in the Clinical Setting

- How should alterations in biomarkers due to treatment be managed (e.g., adrenal insufficiency secondary to systemic treatments)?
- How can allostatic load (AL) be integrated into clinical risk assessment?
- How can we treat elevated allostatic load?

Feasibility Study of Biobehavioral Stress Reduction Intervention in Patients With Triple Negative Breast Cancer (NCT05677802)

STUDY POPULATION



N=40 Age 18+ Untreated TNBC Stages I-III

INTERVENTION

Week 1

- Progressive muscle relaxation
- Responding to cancer crisis (stress, anxiety, depression)
- Physical activity log

Week 2

Coping (seeking disease info)

Weeks 3-5

problem solving (HRSN)

Week 6-7

 assertive communication with healthcare providers

Week 8-10

social support

OUTCOME

Program Acceptability

Allostatic load


Patient Reported Outcomes

- Self reported stress
- depressive symptoms
 - anxiety
 - QoL

Obeng-Gyasi et al, Feasibility Study of Biobehavioral Stress Reduction Intervention in Patients With Triple Negative Breast Cancer (NCT05677802), 2022

Take Away

- Ecosocial theory and weathering describe internalization of socioenvironmental factors.
- Allostatic load is a form of embodiment/weathering.
- High allostatic load is associated with poor tumor characteristics and worse mortality















Acknowledgements

- Barbara Andersen PhD
- William Carson MD
- JC Chen MD
- Mohamed Elsaid PhD
- Demond Handley, MA
- Timothy Pawlik MD, PhD, MBA





• Email: Samilia.obeng-gyasi@osumc.edu

• X: @GyasiSamilia