


## The people who did the work



JTGH Lab 10/2018

#### GENE x ENVIRONMENT



Genetic
Makeup

Common
Environmental
Exposures



#### Parkinson's disease incidence

**RARE** 

COMMON

RARE

## PARK8: LRRK2

- The most common cause of autosomal dominant PD May account for 10% of AD PD and ~3% of sporadic PD
- LRRK2 is a kinase substrates still being defined and verified (Rabs 8 & 10)
- More than 40 point mutations have been identified At least 6 are clearly pathogenic - generally associated with increased kinase activity
- The Gly2019Ser mutation is the most prevalent and is present in 30-40% of PD patients (familial & sporadic) in North African populations & in 13% of Ashkenazi Jews
- Incomplete penetrance
- Role of LRRK2 in idiopathic PD has been unclear







- Growing consensus that S1292 phosphorylation correlates with activity. "...Ser(P)-1292 LRRK2 may be an effective biomarker surrogate for LRRK2 kinase activity"
- 14-3-3 may bind to the inactive form of LRRK2 and 14-3-3 overexpression decreases kinase activity while 14-3-3 dissociation promotes kinase activity
- Can we take advantage of these properties?






#### A multiplex PL assay to assess:

- <u>S1292 phosphorylation state</u>: Amplification allows low Ab concentration (less n.s. binding) and PL eliminates signal from "off-target" Ab binding
- 14-3-3 binding to LRRK2: Allows assessment of endogenous protein:protein interactions in situ
- Validation in <u>CRISPR/Cas9-edited LRRK2<sup>-/-</sup> and LRRK2<sup>G2019S/G2019S</sup> cells
  </u>









Di Maio et al, Sci Transl Med, 2018


















## Is LRRK2 activation reproduced in animal models of PD?



## Rotenone

- Classical high-affinity inhibitor of complex I of the mitochondrial electron transport chain
- A natural product from several plant species

**PIND** 

Pittsburgh Institute for

Neurodegenerative Diseases

- Common pesticide; the "organic" (natural) alternative to synthetic pesticides
- Used to sample fish populations in reservoirs & kill nuisance fish in lakes
- Highly lipophilic; crosses biological membranes easily
   \* independent of transporters



## FAME Study: PD in Agricultural Health Study Tanner, Kamel et al

Nested, case-control study of 52,000 licensed pesticide applicators and 32,000 spouses

Rotenone — Increased Risk of PD:

All OR = 2.3 (95% C.I.: 1.2, 4.3)

Men OR = 2.8 (95% C.I.: 1.4, 5.8)

Model adjusted for age, gender, state, ever smoking, ever pesticide use



Plant derivative (worldwide)


Crop insecticide (mid-1800's - present)

Fish poison (ancient)




Home gardening






Pest Control








The rotenone model











# Redox activation of LRRK2? Effects of H<sub>2</sub>O<sub>2</sub>









#### **Preliminary conclusions:**

- LRRK2 is activated in nigrostriatal neurons in brains of people with iPD
- This is replicated in the rotenone rat model of environmental exposure
- α-Synuclein overexpression also activates LRRK2 in nigrostriatal neurons
- LRRK2 appears to be redox-activated (common to both mitochondrial defects and α-synuclein overexpression)
















TABLE 3: Solvent Exposure Frequencies and Adjusted Pairwise Odds Ratios in PD-Discordant Twins, n = 99Pairs<sup>a</sup>


Solvent	Case <sup>-</sup> / Control <sup>-</sup>	Case <sup>+</sup> / Control <sup>-</sup>	Case <sup>-</sup> / Control <sup>+</sup>	Case <sup>+</sup> / Control <sup>+</sup>	Ever/Never Exposed, OR (95% CI)	p
Toluene	72	11	9	7	1.3 (0.5–3.3)	>0.2
Xylene	88	6	2	3	2.2 (0.4–12)	>0.2
<i>n</i> -Hexane	85	6	7	1,	1.3 (0.4–4.1)	>0.2
CCl <sub>4</sub>	74	14	9	2	2.3 (0.9–6.1)	0.088
PERC	93	5	1	0	10.5 (0.97–113)	0.053
TCE	87	9	2	1,	6.1 (1.2–33)	0.034
TCE or PERC	85	11	2	1	8.9 (1.7–47)	0.010
Any of 6 solvents	51	19	14	15	1.7 (0.8–3.7)	0.16
Any of 4 excluding TCE and PERC	53	18	15	13	1.5 (0.7–3.1)	>0.2

 $^a$ Ever exposure, adjusted for respondent type and smoking.  $CCl_4 = carbon$  tetrachloride; CI = confidence interval; OR = odds ratio; PD = Parkinson disease; PERC = perchloroethylene(tetrachloroethylene); TCE = trichloroethylene.






#### Timecourse of toxicant induced ROS in N27-A (h-αSyn) cells



#### Timecourse of toxicant induced LRRK2 activity in N27-A (h-αSyn) cells



De Miranda et al., unpublished

## Modeling TCE exposure in vivo




#### REQUIREMENTS:

- 1. Produce mitochondrial dysfunction —> oxidative stress
- 2. Recapitulate Parkinsonian nigrostriatal pathology
- 3. Limited off-target toxicity
- 4. Relevant human dose






#### Striatal TH Intensity




\*n=6 control, 10 TCE






#### Stereology






\*n=6 control, 10 TCE

#### TCE exposure induces LRRK2 in dopaminergic neurons








\*n=6 control, 10 TCE



#### pSer129-αSyn










## Role of LRRK2 in endolysosomal function in PD

- 1. Use compartment-selective markers to define status of endolysosomal function in human SNc neurons in iPD
- 2. Determine the extent to which this is replicated in the rotenone model of PD
- Assess whether LRRK2 kinase inhibition ameliorates impaired endolysosomal function and protects SNc neurons






#### Healthy neuron






## Early Endosomes in iPD








## Early Endosomes in the Rotenone Model








## Late Endosomes & Lysosomes in iPD







## Late Endosomes & Lysosomes in the Rotenone Model







## Role of LRRK2 in endolysosomal function in PD

#### In iPD:


- Accumulation of early endosomes
- Depletion of late endosomes & lysosomes (& Cathepsin D)
- Loss of GCase activity and protein levels
- All of these findings are reproduced in the rotenone model

Does LRRK2 kinase inhibition ameliorate impaired endolysosomal function and/or protect SNc neurons of rats treated to rotenone **endpoint**?






# Effects of LRRK2 Inhibition: Target engagement








## Early Endosomes: Effects of LRRK2 Inhibition








## Late Endosomes & Lysosomes: Effects of LRRK2 Inhibition








## Autophagosomes: Effects of LRRK2 Inhibition








## alpha-Synuclein: Effects of LRRK2 Inhibition








## SNC cell loss: Effects of LRRK2 Inhibition
















## LRRK2 Therapeutics in iPD

## Implications:

- LRRK2 plays a central role in iPD, at least in part, by modulating endolysosomal function
- Rather than targeting only the 2-3% of PD patients with LRRK2 mutations, LRRK2 therapeutics may be useful for individuals with iPD, independent of mutations





# Parkinson's disease: At the intersection of genes & environment

#### Conclusions:

- Genes implicated in PD (e.g., α-synuclein & LRRK2) strongly influence the neurotoxicity of environmental toxicants (e.g., rotenone, TCE).
- PD-relevant environmental toxicants (e.g., rotenone, TCE) can "activate" gene products implicated in PD (e.g., pS129-α-synuclein and LRRK2).





## Funding











