

Mechanisms Associated with Dietary Energy Balance Effects on Tumor Development: Alterations in Growth Factor and Energy Sensing Pathways

John DiGiovanni, Ph.D.

**Professor and Coulter R. Sublett Endowed Chair
Division of Pharmacology and Toxicology
College of Pharmacy and Department of Nutritional
Sciences, College of Natural Sciences
The University of Texas at Austin**

Presentation Outline


- **Part I**

- Dietary energy balance modulates skin tumor promotion via altered growth factor signaling (e.g., Akt/mTOR) and proliferation during tumor promotion.
- Possible role of altered IGF-1R/EGFR crosstalk in dietary energy balance effects on tumor promotion.
- Targeting mTORC1 as a strategy for inhibition of tumor promotion mimicking the effects of CR.

- **Part II**

- Dietary energy balance modulates prostate cancer progression in Hi-Myc mouse model
- Dietary energy balance effects growth factor signaling pathways in prostate
- Possible role of inflammation and inflammatory signaling in prostate cancer progression in Hi-Myc Mice

Potential Mechanisms Underlying the Link Between Energy Balance and Cancer



Dietary Energy Balance Modulation Alters Steady State Growth Factor Signaling Pathways in Epithelial Tissues of Normal Mice


Epidermis
Liver
Prostate

Caloric Density of Diets Used to Examine the Effect of Dietary Energy Balance on Growth Factor Signaling in Epithelial Tissues





<i>Experimental diet</i>	<i>Experimental Phenotype</i>	<i>Protein (kcal%)</i>	<i>Carbohydrate (kcal%)</i>	<i>Fat (kcal%)</i>
<i>30% CR¹</i>	Lean	29	57	14
<i>15% CR²</i>	Normal	24	65	12
<i>AIN76A³</i>	Overweight	20	70	10
<i>DIO</i>	Obese	20	20	60

Effect of Dietary Energy Balance Modulation on Weight Distribution and Serum Hormone Levels in Male Mice



FVB Mice



C57BL/6 Mice





Effect of Dietary Energy Balance Modulation on Growth Factor Signaling in Epithelial Tissues of Untreated Mice



Dietary Energy Balance Modulation and Two-Stage Skin Carcinogenesis

- Calorie restriction (CR) significantly reduces tumor incidence, multiplicity, and papilloma size during two-stage skin carcinogenesis (Boutwell, 1964; Birt et al, 1991; Birt et al, 1993).
- Inhibitory effects of CR are observed primarily during tumor promotion (Birt et al, 1991).
- Potential mechanisms of inhibition:
 - § 40% CR inhibits ERK, leading to decreased proliferation (Liu et al, 2001);
 - § Levels of corticosterone may mediate inhibitory effects of 40%CR (Stewart et al, 2005);
 - § Reduced PI3K and Ras signaling following treatment with TPA (Xie et al, 2007);
 - § Altered growth factor signaling via reduced circulating IGF-1 (Moore et al, CPR 2008; Cancer Res., 2008)
- Effect of positive energy balance is less clear.

Dietary Energy Balance Modulation of Skin Tumor Promotion by TPA in ICR Mice



Moore et al, unpublished data

Dietary Energy Balance Does Not Affect the Rate of Malignant Conversion of Papillomas to SCCs

Experimental Diet	Total Number of Mice	Average Papillomas Per Mouse ¹	Carcinoma Incidence (%)	Carcinomas per Mouse	Conversion Ratio ²
60kcal% fat	25	8.86	96.0	2.28 ^a	0.26
10kcal% fat	27	8.20	92.3	1.58	0.20
15% CR	29	6.17 ^a	69.0	1.59	0.26
30% CR	26	4.27 ^a	57.7	0.96 ^a	0.23




¹Data taken at 27 weeks of promotion with 3.4 nmol TPA after which the papilloma response had reached a plateau

²Ratio of the average number of SCCs at 50 weeks to average number of papillomas at 27 weeks





^aSignificantly different from all groups, Wilcoxon Rank Sum (p<0.05)

Moore et al, unpublished data

Dietary Energy Balance Modulation Alters Epidermal Proliferation Following TPA Treatment




Weight Gain and Selected Serum Protein Profiles of Female ICR Mice in the Two-Stage Skin Carcinogenesis Study



^{a,b} Significantly different from values with the same lettering Wilcoxon Rank Sum ($p < 0.05$)

Moore et al, unpublished data

Summary of Dietary Energy Balance Effects on Growth Factor Signaling and Cell Cycle Related Proteins in Epidermis During Tumor Promotion




Moore et al, CAPR, 2008 and unpublished data

Reduced Circulating IGF-1 Inhibits Two-Stage Skin Carcinogenesis

LID Mice



- Circulating IGF-1 is produced primarily by the liver
- Cre/loxP system was utilized to create a targeted deletion of liver specific IGF-1 production
- Results in a 75% reduction in circulating IGF-1
- Targeted deletion had no effect on post-natal growth
- Liu et. al. *Proc Soc Biol Exp Med*, 2000; Yakar et. al. *PNAS*, 1999



TPA 13.6, Wt


TPA 13.6, LID

Reduced Circulating IGF-1 Decreases Epidermal Proliferation Both in the Absence and Presence of TPA Treatment




*Significantly different ($p < .05$) from values obtained from wild-type mice


Reduced Circulating IGF-1 Decreases Activation of EGFR, IGF-1R and Akt/mTOR Signaling Induced by TPA






Mechanisms for IGF-1R/EGFR Cross-Talk



Targeting mTORC1 Signaling





Rapamycin is a Highly Potent Inhibitor of Skin Tumor Promotion by TPA




* denotes significance. Differences in the average number of papillomas per mouse at 25 weeks between the TPA control group and corresponding Rapamycin treated groups for each dose were statistically significant ($P < 0.05$, Mann-Whitney U Test).

* Significantly different from 6.8nmol TPA ($P < 0.05$, Chi-square test).

Rapamycin Treatment Inhibits TPA-Induced Epidermal Hyperproliferation





Topical Rapamycin Inhibits TPA-Induced Epidermal mTORC1 Signaling Primarily Through the p70S6K Downstream Signaling Pathway



Rapamycin Potently Inhibits Skin Tumor Promotion by TPA in Overweight and Obese Mice





Overweight






* Indicates significantly different from 6.8 nmol TPA (p <.05)

Obese




Checkley et al, unpublished

Metformin Given in the Drinking Water Inhibits Skin Tumor Promotion by TPA in Overweight Mice



Checkley et al, unpublished

mTORC1 Signaling Pathway as a Potential Target for Prevention of Obesity Related Cancer Promotion



Hi-Myc Mouse Model of Prostate Cancer

- Overexpression of c-myc in ARR2PB-myc-PAI transgenic mice resulted in complete penetrance of PIN as early as 2 to 4 weeks of age.
- Progression to locally invasive adenocarcinomas occurs within ages 3 to 6 months for Hi-myc mice and within 10 to 12 months in mice expressing low c-myc levels.
- Ventral Prostate, Dorsolateral Prostate, and Anterior Prostate are the target of c-Myc overexpression.
- Metastases are not present in these transgenic mice.
- This mouse model was the first nonSV40 model to develop advanced adenocarcinoma without NE features.

Effect of Dietary Energy Balance Manipulation on Body Mass, Body Fat Content and Circulating Growth Factors/Adipokines in Hi-Myc Mice.

	30% CR	Overweight control	DIO
Initial mass (g)	25.9 ± 0.9	24.1 ± 0.96	22.3 ± 0.88
Final mass (g) ^b	23.9 ± 1.2	40.1 ± 1.2	44.9 ± 1.6
Body fat (%) ^b	21.1 ± 0.84	38.0 ± 0.87	41.2 ± 1.2
Feed consumption (g/wk) ^b	21.2 ± 0.04	29.8 ± 0.43	28.1 ± 0.8
IGF-1 (ng/mL) ^b	383.7 ± 47.1	550 ± 56.8	893. ± 90.9
Insulin (pg/mL) ^b	2624.6 ± 235.9	3934.2 ± 557.9	5682.9 ± 557.5
Leptin (pg/mL) ^c	801.6 ± 293.9	10385.9 ± 960.3	10392.1 ± 608.9
Adiponectin (ng/mL) ^c	12205.7 ± 898.5	9417.3 ± 666.9	9654.7 ± 568.2
Resistin (pg/mL) ^b	1516.1 ± 84.7	2003.7 ± 213.4	2538.5 ± 167.3

^a N=12 mice/group; values are means ± SEM. Serum analyses were performed at study's end.

^b Value for each diet group differed significantly (p<0.05, Wilcoxon rank sum) from that in the other groups for this parameter.

^c Value differed significantly (p<0.05, Wilcoxon rank sum) in the 30% CR group for this parameter relative to the other two groups.







Characterization of Representative Lesions in Ventral Prostate HI-Myc Mice on the Different Diets at 6 Months

30% CR

Overweight Control


Obese

6 months




Dietary Energy Balance Effects Progression of Prostate Cancer in Hi-Myc Mice on At Both 3 and 6 Months


3 Months



6 Months




Dietary Energy Balance Manipulation Modulates Growth Factor Signaling Pathways in Prostate of Male C57BL/6 and FVB/N Mice










Moore et al, Cancer Prev Res, 2008 and unpublished




Dietary Energy Balance Effects Growth Factor Signaling in Ventral Prostate of Hi-Myc Mice



IHC Staining for Ki67, Cyclin D1, CD31 and p-NFkB IN Ventral Prostate of Hi-Myc Mice




Dramatic Increases are Seen in Stromal Inflammatory Cell Infiltration in Ventral Prostate of Hi-Myc Mice on an Obesity Inducing Diet




Perilipin Staining of Adipocytes in Ventral Prostate of Hi-Myc Mice on Different Diets at 6 Months


30% CR




Overweight Control




Obese



Dietary Energy Balance Dramatically Effects Expression of Inflammatory Cytokines and VEGF Family Members




IL-23 levels are dramatically upregulated




The IL-23 Receptor is Highly Upregulated in Ventral Prostate of Hi-Myc Mice on an Obesity Inducing Diet at 6 Months


30% CR




Overweight Control



Obese



Obesity Increases Prostate Cancer Progression Through Multiple Mechanisms



From Kruijsdijk et al, CEBP 2009

Acknowledgements

DiGiovanni Lab

Tricia Moore, Ph.D.
Allyson Checkley
Okkyung Rho, PhD
Kaoru Kiguchi, MD
Jorge Blando, DVM
Achinto Saha, Ph.D.
Steve Carbajal
Linda Beltran

Collaborators

Steve Hursting, PhD (UT Austin)

Research Supported by NIH Grants CA37111
CA129409 and CA140388. TM was supported by NIEHS
Training Grant ES007247