
# Offshore Wind Farm Impacts on Pacific Upwelling, Nutrients, and Productivity





#### **Project Team**

Mike Jacox



Jerome Fiechter

Ludivine Conte



Grace Chang



Kaus Raghukumar



Chris Chartrand





















Farr et al. (2021)






Do offshore wind turbines have the potential to impact coastal upwelling and consequently nutrients and productivity?

Project funded by California Energy Commission (2020-2023)



Modeled wind-farm induced changes in the North Sea



### **Project Goals and Objectives**

1. Evaluate the potential effects of California OSW farms on circulation (i.e., upwelling) and biogeochemistry (nutrients, productivity) off the U.S. west coast.


2. Compare simulated biogeochemical impacts of OSW farms with impacts projected due to climate change in coming decades.

### **Project Goals and Objectives**

1. Evaluate the potential effects of California OSW farms on circulation (i.e., upwelling) and biogeochemistry (nutrients, productivity) off the U.S. west coast.

2. Compare simulated biogeochemical impacts of OSW farms with impacts projected due to climate change in coming decades.

### **Modeling Configuration**



### **Atmospheric Modeling**


Weather Research and Forecasting (WRF) model with Wind Farm Parameterization

Morro Bay, Humboldt, Coos Bay, Brookings wind energy areas

15MW and 20MW turbine build-outs


3km resolution for full domain

1995-2019




### **Atmospheric Modeling - Wind Farms**

#### PRIOR STUDY



10 MW Turbines; 9D spacing

#### THIS STUDY



15 MW or 20 MW Turbines; 10x4D elliptical spacing

### Ocean Modeling

3km resolution for full domain

1995-2019

Atmospheric forcing w/ and w/out turbines


Physics (ROMS) + Biogeochemistry (NEMUCSC)

**Nutrients** 


Oxygen

Two phytoplankton groups

Three zooplankton groups





### Wind Speed




972 x **15 MW** turbines = **14.6 GW** total 595 x **20 MW** turbines = **11.9 GW** total


Upwelling Season (March - July)
Average of 15MW and 20MW simulations






Upwelling Season (March - July)
Average of 15MW and 20MW simulations






### **Phytoplankton Biomass**


# Upwelling Season (March - July) Average of 15MW and 20MW simulations







### **Zooplankton Biomass**


# Upwelling Season (March - July) Average of 15MW and 20MW simulations








Plankton changes are offset from upwelling changes due to temporal lags + advection





Changes in upwelled nitrate



Particles released in area of decreased nitrate supply










Particles released in area of decreased nitrate supply









# Particles released in area of increased nitrate supply




Particles released in area of decreased nitrate supply









Particles released in area of increased nitrate supply







Changes in phytoplankton



Changes in zooplankton






### **Project Goals and Objectives**

1. Evaluate the potential effects of California OSW farms on circulation (i.e., upwelling) and biogeochemistry (nutrients, productivity) off the U.S. west coast.

2. Compare simulated biogeochemical impacts of OSW farms with impacts projected due to climate change in coming decades.

### Offshore Wind Impacts in the Context of Climate Change


Projected end-of-century changes under high emission scenario



Jacox et al. (2024)

### Offshore Wind Impacts in the Context of Climate Change

Projected end-of-century changes under high emission scenario



### Offshore Wind Impacts in the Context of Climate Change

Wind farm impacts are patchy mosaics of positive and negative. Neither natural variability nor climate change induce similar patterns in upwelling.

Impacts of northern areas (Humboldt, Coos Bay, Brookings) are harder to discern, likely because they are smaller and interact with each other.

Nearshore, wind farm impacts on phytoplankton are smaller than projected climate change impacts (under high emissions scenario). Offshore, they can be comparable.

Presumably, more offshore wind development accompanies lower emissions scenarios. If so, the relative impact of offshore wind will increase as emissions decrease.