


A little about myself...

Research Scientist; Volcanologist; PI of Moon Diver and MoonDrop concepts





Moon Diver was a 2019 Discovery Concept to explore a pit in Mare Tranquillitatis


Plan for the Talk


- Pit Basics: Types and Distribution
- Caves, Pits, and Lunar Hazards: Radiation, Thermal Swings, Micrometeorites, Dust, Vacuum
- Pros and Cons of Specific Sites
- What Can Humans Do Best? (and how can robots help?)
- Access and Stability
- References

Pit Basics: Impact Melt Pits vs Deep Pits

Impact Melt Pits (257)

Mare (15) and Highland (6) Pits






Important Points:


Wagner and Robinson, 2014


Impact Melt pits are numerous but small (median diameter: 15 m). Mare pits are few but large. 256 total pits measured [by Wagner et al., 2017], 178-194 are at least large enough to host a 3-person habitat. 30-40% of measured pits have some kind of "ramp". Only a handful of pits are likely to have caves (mostly mare pits). Impact melt pits are much less scientifically interesting than mare pits. Consult the LRO Pit Atlas for more details!! https://lroc.im-ldi.com/atlases/pits

Pit Distribution



Pit Distribution





Important Points:

- There are no confirmed pits in the polar regions (>84° latitude)
- For Mare Pits, it is because there is no mare!
- For Impact Melt Pits, it is hard to confirm them
- Even Pascal Lee's northern pits are outside the polar region (~73°N)
- Pits are not expected to contain water ice (even if they are at high latitudes) (Wilcoski et al., 2023); however, water left by humans could persist for some time.




The Big 5







Presence of a Ramp Presence of a Cave



Science

Regolith Examining the regolith/bedrock interface (in-situ)

Physical Volcanism Determining how magmas are transported to the surface on the Moon (in-situ and sample return for absolute chronology)

Petrology Determining the structure of the Moon's interior (in-situ or sample return)


History of the Magnetic Field (in-situ or sample return)

Solar History (shallow paleoregolith sample return)


Galactic History (deeper paleoregolith sample return)







Thick or thin?





LUNAR HAZARDS

RADIATION

MICRO-METEORITES THERMAL SWINGS

DUST




MICRO-METEORITES

THERMAL SWINGS

DUST

VACUUM

- Radiation is a major limiting factor for long-term stays on the Moon. The annual radiation dose on the Moon is ~11% of NASA's career dose limit for an astronaut (and ~25% of ESA's), before accounting for solar flares [1,2,3].
- There are two major sources of radiation on the Moon: Solar radiation and cosmic ray nucleii.



Modified by Wagner et al. from De Angelis, G. et al. (2007)



MICRO-METEORITES THERMAL SWINGS

DUST


VACUUM



Regolith Shielding


- Can build anywhere
- Full protection requires moving 4+ m of regolith into place
- Complex engineering and materials simulation needed

- Difficult to expand



Caves

- Limited to places with caves
- Extremely good protection
- Easy to use inflatable habitats
- Can't conclusively identify cave entrances from orbit



Pits


- More common than cave access
- Moderate protection
- Likely easier access than caves
- Often limited interior space
- Limited line-of-sight to Earth for communication


MICRO-METEORITES

THERMAL SWINGS

DUST

- Of the 256 measured pits, 178-194 are large enough to hold a three-astronaut habitat module (depending on model shape). 40-49 of these pits reduce radiation by 83% or more, corresponding to an annual dosage of less than 50 mSv (the limit for U.S. radiation workers [2]).
- Some pits (~43%) have an additional feature that would be useful for astronauts: a collapsed wall that provides possible access by driving, rather than building an elevator.
- Unfortunately, pits with these collapses are rarely narrow and deep enough to provide good protection. Of the pits with <50 mSv/yr exposure, only ~8 have possible entrance ramps.







DUST

- The Grun Lunar flux model estimates that a micrometeoroid with a mass equal to one microgram will impact the lunar surface twice per square meter per year, causing damage ranging from catastrophic failure to partial penetration and pitting, local deformation, and material degradation. [Allende et al., 2020]
- Protection is also needed from nearby rocket blasts









MICRO-METEORITES


THERMAL SWINGS

DUST




Horvath et al., 2022





MICRO-METEORITES THERMAL SWINGS


DUST



MICRO-METEORITES THERMAL SWINGS


DUST


VACUUM



Horvath, Hayne, and Paige (2022). Thermal and Illumination Environments of Lunar Pits and Caves: Models and Observations from the Diviner Lunar Radiometer experiment









MICRO-METEORITES

THERMAL SWINGS

DUST

- The floor of lunar pits are likely to have slightly more dust than their surroundings (because dust can fall in but it cannot get out)
- There will be little to no dust actually inside the tubes.



Kazumura Cave, Hawai'i (65 km long)

MICRO-METEORITES


THERMAL SWINGS

DUST

VACUUM

Can lava tubes be hermetically sealed?

No; but....



Permeability- Instead, permeability in lavas comes mostly from a combination of boundaries between lavas (horizontal) and jointing from cooling (vertical). Dense lava cores have very low permeability.


Cracks On Earth, cracks often form during cooling and then widen later due to infiltration by water, earthquakes, and tectonic stresses.




Cueva de los Verdes, Laura Kerber

Lacus Mortis Pit









Low iron, low titanium

Lacus Mortis




The devil is in the details on ramps: we need higher resolution imagery!

Mare Ingenii Pit



Lat -35.949 Lon 166.056







Marius Hills Pit



Lat **14.0917** Lon 303.2299







Tranquillitatis Pit



Lat **8.3355**. Lon **33.222**




Located on near-side
Best quality imagery
Can see the Earth from the bottom
Has a cave (based on visible data,
thermal anomaly and Mini-RF)
Inner diameter: 100 x 88 m
Depth: 105 m
Close to intersection of two lava units
High titanium lavas (6-8% TiO₂)
High iron lavas (17.5-19.5% FeO)




Attempted Ways to Confirm a Cave

- Looking with LRO off-nadir (at least 20 m of overhang at Tranquillitatis; at least 14 m at Marius Hills; at least 10-20 on the W and S sides of Ingenii; Wagner and Robinson, 2014).
- Thermal anomaly (confirmed at Tranquillitatis and Ingenii) with Diviner (Horvath et al., 2022)
- Radar (SELENE radar and Mini-RF Kaku et al.; Carrer et al., 2024)
- Gravity (Chappaz et al., 2017)




What Can Humans Do? Step 1



Surface-based exploration:


- -Geophones/seismic
- -Mortars/thumpers
- -Ground penetrating radar
- -Magnetic surveys
- -Gravimeters



Measure subsurface void

Compare geophysical data of lava and regolith layers and compare with ground-truth in outcrop

What Can Humans Do? Step 2




Funnel-based Exploration:

- -Tethered drilling and exploration
- -Examination of the regolith-to-bedrock transition



30-40 m across (~100 ft)









Pit-Based Exploration

Humans could install a hovering camera, similar to the SkyCam and SpiderCams used by the NFL. Humans are good at setting anchors!

Humans could also fly a hopper in real time to explore and map the cave interior.




What Can Humans Do? Step 4





What Can Humans Do? Step 5



But Laura: I don't want to carry all of my stuff down there!







People rappel into these pits on Earth fairly frequently. Things we worry about when we do that:

- Material falling down from above
- Overhangs
- Standing on unstable ledges


Advantages:

Humans are very good at setting anchors! Humans are very good at drilling into walls! Humans are very good at rappelling!




PIT FLOOR

The floor of pits are covered in giant boulders. It is scary, but humans are also actually pretty good at this.







LAVA TUBE FLOOR




Lava River Cave Arizon:





Photo: Lanny El Malpais, NM Wagner











Pretty Stable, All Things Considered

Types of Instability:

Blocks- Mitigation: Inspection, Tilt-meters

Gravel and Sand- Mitigation: Inspection, Mortar, and Resin

Plates – Mitigation: Inspection, Bolts if needed

Span—Mitigation: Inspection, Modeling

Inspections at major tourist caves are mostly done after major events such as earthquakes, floods, and volcanic eruptions.

These instabilities are less likely to be a problem on the Moon in the absence of water and chemical weathering. For example, the fall of gravel and sand is mostly due to chemical weathering of the rocks due to water flowing through cracks.



Mount Fuji Lava Tubes (Kerber)




Hengifoss, Iceland


STABILITY

Things to observe and measure:

- Are the lavas thick or thin? (tens of cm vs meters vs decameters)— thick lava flows will be more stable than thin ones
- Are the lavas porous or dense?
- Are the lavas a'a or paehoehoe? A'a is thicker and denser, except at the boundaries between lavas. Pahoehoe is what forms tubes.
- Are there granular layers between lavas? (paleoregolith; clinkers)
- Are the original lava tube textures still present? (lavacicles, flows on the floor, glazed walls)
- What is the density (and orientation) of cracks?
- How much break-down is there? (fallen ceiling blocks, blocks covering the floor)







Signorelli et al., 2019

Conclusions

- Lunar pits are a desirable non-polar landing site for planetary science, astrophysics, heliophysics, biological and physical sciences, and applied sciences.
- There are many possible pits that can be visited, but the most attractive scientifically and for a long-term base are the large mare pits, especially those with caves. There is no known pit with both a ramp and a "known" cave.
- More information is needed before we can commit to building a habitat in a lunar pit, including internal geometry, amount of breakdown, lava types, thermal and radiation environment measurements.
- Humans are typically very good at the types of activities that would be needed (geophysical surveys, setting anchors, rappelling, rock-hopping).
- It would be useful to have some precursor information about the detailed geometry of the pit to be visited, if possible.

Applied Sciences

Goal	Measurements	Mission Requirement
Determine size and shape of cave	Local seismic, magnetic, gravimeter, radar	Surface is ok
Determine size and shape of cave (high quality)	Lidar	Peek into cave (descend to overhang)
Determine stability of cave	Images of cave to see if there is breakdown vs pristine lava-tube structures, assessment of thick or thin lava layers	Peek into cave (descend to overhang)
Determine entrance geometry	Images of cave or lidar	Peek into cave; further entrance may be needed
Determine density/porosity of lavas	Close-up examination of lavas in wall	Access to wall
Identify paleoregolith (for stability reasons)	Images of walls	Top of funnel
Understand the regolith/bedrock interface (for ISRU)	Geophysical measurements; direct imaging	Surface; top of funnel

Goal	Measurements	Mission Requirement
Determine thickness and morphology of lava	Images of walls	Close-up flyover is ok; access to walls is preferred
Determine rheology of lavas	Bulk elemental composition; bubble and crystal distributions	Access to walls, can be done in-situ, samples would be better
Understand the regolith/bedrock interface		Peek into cave (descend to overhang)
Determine entrance geometry	Images of cave or lidar	Peek into cave; further entrance may be needed
Determine density/porosity of lavas	Close-up examination of lavas in wall	Access to wall
Identify paleoregolith	Images of walls	Top of funnel






More References

- Habitability Potential of Lunar Pit Craters: Marius hills, Mare Tranquillitatis, Lacus Mortis, and Mare Ingenii Pit https://www.hou.usra.edu/meetings/lpsc2023/pdf/2380.pdf
 Wagner, R. V., and M. S. Robinson. "Lunar pit morphology: Implications for exploration." *Journal of Geophysical Research: Planets* 127.8 (2022): e2022JE007328.
- Overview of the CHILL-ICE 2021 Science Experiments and Research Campaign (field experiment deploying an inflatable habitat in a lava tube) https://spi.science.org/doi/full/10.34133/2022/9760968
- A comprehensive review of lunar lava tube base construction and field research on a potential Earth test site https://www.sciencedirect.com/science/article/pii/S2095268624000752#b0280
- Horvath, T., Hayne, P. O., & Paige, D. A. (2022). Thermal and illumination environments of lunar pits and caves: models and observations from the diviner lunar radiometer experiment. *Geophysical Research Letters*, 49(14), e2022GL099710. https://agupubs.onlinelibrary.wiley.com/doi/full/10.1029/2022GL099710
- Halliday, W. R. (1998). "Pit Craters", lava tubes, and open vertical volcanic conduits in Hawaii: a problem in terminology. International Journal of Speleology, 27(1), 12. https://digitalcommons.usf.edu/cgi/viewcontent.cgi?article=1278&context=ijs
- Signorelli, S., Carmona, F. J., Cabrera, M. P., & Zafrilla, S. (2019, August). The Jameos del Agua cave (Lanzarote, Canary Islands): Some morphological and geological features of a spectacular lava tube. In *Volcanic Rocks: Proceedings of ISRM Workshop W2, Ponta Delgada, Azores, Portugal, 14-15 July, 2007*. CRC Press. <a href="https://books.google.com/books?hl=en&lr=&id=S5DUDwAAQBAJ&oi=fnd&pg=PA45&dq=jameos+del+agua&ots=4jQIF-yO3D&sig=VVwltJ4ZdZS1SXKP43cOo7FiDAg#v=onepage&q=jameos%20del%20agua&f=false
- Hernández-Gutiérrez, L. E., Santamarta, J. C., Pacheco, L., Martín-González, E., Hernández-Martín, H., Xifré, R., & Calderón-Guerrero, C. (2025). Tools for Managing the Integrity of Tourist Volcanic Caves in the Canary Islands Due to Instability Problems. *Geosciences*, 15(7), 236. https://www.mdpi.com/2076-3263/15/7/236



Science

• Physical Volcanism: Determining how magmas are transported to the surface on the Moon










Is it local?

Is it representative?



Neal et al., 2015

Is it an igneous rock?



Neal et al., 2015




Where in the lava does it come from?




Science

 Petrology: Piecing together the structure of the lunar mantle and crust with petrology

Is it a liquid composition?
Is it a primary composition?




KNOWLEDGE OF THE LUNAR INTERIOR!




Science

 Regolith: Exploring the regolith/bedrock interface and process of regolith formation

 Comparing the surface regolith to the underlying layers to determine its representativeness





Science

Lunar Magnetism: Did the lunar crust have a late dynamo?

Paleoregolith: What was the density of the ancient solar wind?

