
# Mental Health and Aging in Native Communities: Adversity, Innovative Solutions, Survivance




Spero M. Manson, PhD Distinguished Professor of Public Health and Psychiatry Director, Centers for American Indian and Alaska Native Health Colorado School of Public Health University of Colorado Anschutz Medical Campus

## The Demographic Transition in Native Communities



 1970 Life expectancy of American Indian male was 47.7 years
2019 Life expectancy of

American Indian male was 71.8 years



## The Epidemiologic Transition in Native Communities

#### Epidemiologic Transition

	Stage 1	Stage 2	Stage 3	Stage 4	Stage 5
_	Infectious and parasitic diseases Accidents and animal attacks "Natural checks" on population	Receding pandemics Sanitation, nutrition, medicine lead to lower CDR	Degenerative and man- made diseases Heart disease, cancer, diabetes, obesity	Delayed degenerative diseases Extend life expectancy due to medical advances	Potential resurgence of infectious diseases due to globalization
			CHRONIC DISEASES		


Epidemiological transition represents a shift in the disease pattern of a population as mortality falls during the demographic transition: acute, infectious diseases are reduced, while chronic, degenerative diseases increase in prominence, causing a gradual shift in the age pattern of mortality from younger to older ages.






## **Competing Risk and Survivor Resilience**

### Suicide Rate by Age for American Indian/Alaska Native Compared to United States (Average 2000–2013)



#### OPIOID Overdose Deaths among Native Americans by Sex & Age, U.S. 2014-2016



The opioid overdose death rate among Native American males significantly exceeds the rate among Native American females (10.0 per 100,000 vs. 7.0 per 100,000). Opioid overdose deaths are significantly more common among Native Americans between the ages of 25-64.

SOURCE: Centers for Disease Control and Prevention, National Center for Health Statistics. Multiple Cause of Death 1999-2016 on CDC WONDER Online Database

## **Depression and Post-Traumatic Stress Disorder**

- Significantly higher rates of self-reported depressive symptoms among older American Indians than non-Hispanic Whites associated with ...
  - Greater prevalence of chronic health problems such as diabetes and cardiovascular disease
  - Significant limitations in activities of daily living
  - Isolation and decreased ability to participate in social and cultural events
  - Elder abuse and neglect
- PTSD 2-3 times more prevalent among older American Indians and Alaska Natives than non-Hispanic Whites, associated with greater ...
  - Frequency of adverse childhood events
  - Exposure to interpersonal violence



## Mental Illness, Substance Abuse, and Stigma

Stigmatizing Condition* (n=3,043)	Mean	Rank Order
HIV/AIDS	4.9	1
Mentally ill	4.6	2
Attempted suicide	4.3	3
Sexually assaulted	3.9	4
Homeless	3.6	5
Alcoholic	3.0	6
Obese	2.5	7
Divorced	2.2	8
Unemployed	2.0	9


\*\*"How ashamed or embarrassed would you feel if it became known to others that you had or were .... ?" 1= little or not at all; 5=most possible

## Screening, Brief Intervention and Referral to Treatment: Alaska Medical Center, Anchorage, AK



- SBIRT is well-established, evidence-based practice to detect and manage behavioral health problems in primary care
- Medical home for 65,000 Alaska Native/American Indians living in southcentral Alaska
- 36 primary care teams treat empaneled patients; 1 masters-level clinician, known as Behavioral Health Consultant, co-located with each team
- Initial screening for behavioral health problems alcohol, substance abuse, depression, trauma, and suicide – upon presentation for appointment
- Began in 2001; screened over 58,000 unique patients; 27% deemed high risk and followed up; screening expanded over time to embrace wide range of conditions and ages; referred to a continuum of behavioral health care

## **Rural Native Veterans and Telepsychiatry**



### Survivance

